



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> C12N 5/08, C12Q 1/02, A61K 35/14, 38/19	<b>A1</b>	<b>(11) International Publication Number:</b> WO 97/29182 <b>(43) International Publication Date:</b> 14 August 1997 (14.08.97)
<b>(21) International Application Number:</b> PCT/US97/02110 <b>(22) International Filing Date:</b> 10 February 1997 (10.02.97)  <b>(30) Priority Data:</b> 08/600,483 12 February 1996 (12.02.96) US  <b>(71) Applicant:</b> THE ROCKEFELLER UNIVERSITY [US/US]; 1230 York Avenue, New York, NY 10021 (US).  <b>(72) Inventors:</b> STEINMAN, Ralph, M.; 62 North Avenue, West- port, CT 06889 (US). BHARDWAJ, Nina; 114 Upper Mountain Avenue, Montclair, NJ 07042 (US). SCHULER, Gerald; Buckenhoferstrasse 11, D-91080 Spardorf (US).  <b>(74) Agent:</b> SONNENFELD, Kenneth, H.; Morgan & Finnegan LLP., 345 Park Avenue, New York, NY 10154 (US).		<b>(81) Designated States:</b> AU, CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> METHOD AND COMPOSITIONS FOR OBTAINING MATURE DENDRITIC CELLS  <b>(57) Abstract</b>  We describe an improved method for generating sizable numbers of mature dendritic cells from nonproliferating progenitors in human blood. The first step or "priming" phase is a culture of T cell depleted mononuclear cells in medium supplemented with GM-CSF and IL-4 to produce immature dendritic cells. The second step or "differentiation" phase requires the exposure to dendritic cell maturation factor such as monocyte conditioned medium. Using this two-step approach, substantial yields are obtained. The dendritic cells derived from this method have all the features of mature cells. They include a stellate cell shape, nonadherence to plastic, and very strong T cell stimulatory activity. The mature dendritic cells produced according to this invention are useful for activating T cells.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

**METHOD AND COMPOSITIONS FOR OBTAINING  
MATURE DENDRITIC CELLS**

This invention was made with United States Government support under NIH grants AR-39552, AR-42557 and AI-24775. The United States Government has certain rights in this invention.

**TECHNICAL FIELD OF THE INVENTION**

This invention relates to methods and compositions useful for activating cells of the immune system. The methods and compositions provided by this invention are useful for causing the maturation of non-proliferating immature dendritic cells to mature dendritic cells capable of processing and presenting antigen. This invention also relates to culture mediums which promote maturation of immature dendritic cells to mature dendritic cells. In addition, this invention relates to assays useful for detecting the presence in a test substance of a dendritic cell maturation factor.

**BACKGROUND OF THE INVENTION**

Dendritic cells are specialized antigen presenting cells, critical for eliciting T cell mediated immune responses (Steinman, 1991; Caux et al. 1995a; Hart and McKenzie, 1990; Austyn, 1987). These specialized antigen presenting cells elicit both CD4+ helper cells (Inaba et al. 1990a; Inaba et al. 1990b; Crowley et al. 1990) and CD8+ killer cells (Porgador and Gilboa, 1995; Mayordomo et al. 1995; Zitvogel et al. 1995) in vivo.

Because of the potent activity of dendritic cells to activate T cells, the art, e.g. Flanagan et al., Env. J. Immunol., 24:605-610, 1994, has accepted the characterization of dendritic cells as "nature's

-2 -

adjuvant". It is therefore desirable to be able to use dendritic cells to process and present protein antigens as a means of modulating an individual's immune response, and in particular to activate an individual's T cells in connection with the treatment or prevention of disease.

5 The use of primed dendritic cells to activate cytotoxic T cells has been reported. For example, Paglia et al., recently reported that murine dendritic cells cultured from bone marrow precursors and exposed to antigen in vitro provide effective resistance to challenge with live tumor cells. Paglia et al., "Murine Dendritic Cells Loaded In Vitro With Soluble Protein Prime Cytotoxic T Lymphocytes Against Tumor Antigen In Vivop", J. Exp. Med., 183:317-322 (1996).

15 The in vitro observations and the murine results have recently been extended to the treatment of humans with B cell lymphoma using mature dendritic cells which were primed in vitro with tumor antigens. Hsu et al., "Vaccination of Patients With B-cell Lymphoma Using Autologous Antigen-Pulsed Dendritic Cells", Nature Med., 2:52-58 (1996). According to this report, the antigen used to prime the dendritic cells was idiotypic antibody obtained from hybridomas made from the fusion of lymph node tumor cells with a mouse cell line. All of the four patients involved in this study exhibited significant antitumor idiotype PBMC proliferative responses. Positive clinical responses were also observed including one patient who experienced complete tumor regression. Hsu et al., however, express uncertainty whether dendritic cells obtained from expansion of dendritic cell cultures in the presence of GM-CSF, TNF- $\alpha$  or IL-4 would function equivalently in their ability to process and present antigen or to stimulate cellular immune responses as the freshly isolated cells.

35

- 3 -

Prior studies have identified proliferating dendritic cell progenitors within the small CD34+ subfraction of cells in human blood (Inaba et al. 1992; Caux et al. 1994; Caux et al. 1992; Caux et al. 1995). Methods have been developed for expanding these proliferating cells in culture to obtain sufficient numbers of cells to be useful for priming with antigen and administering to an individual to activate an immune response. Steinman et al. International patent application WO 93/20185. These dendritic cells can be stimulated with cytokines, particularly GM-CSF and, optionally TNF $\alpha$  or other cytokines, to develop into potent dendritic cells over 1-2 weeks in culture (Caux et al. 1992; Inaba et al. (1992)). Although useful dendritic cells can be produced from the proliferating precursors, it is desirable to obtain alternative methods of obtaining suitable numbers of mature dendritic cells for therapeutic purposes where proliferating progenitors are infrequent.

The removal of monocytes and lymphocytes from human blood has uncovered a small population of nonproliferating progenitors that require cytokines to develop into typical dendritic cells. These progenitor cells exist at a concentration of at most about  $10^6$  cells per 450-500 ml of blood (O'Doherty et al. 1994; O'Doherty et al. 1993; Thomas et al. 1993). More recently, the combination of GM-CSF and IL-4 has been shown to facilitate the generation of significantly larger numbers of dendritic cells from adherent blood mononuclear fractions, about  $3-8 \times 10^6$  per 40 ml of blood (Romani et al. 1994; Sallusto and Lanzavecchia, 1994 both of which are incorporated herein by reference). However, we have now determined that when the cytokines are removed, the cells revert to an adherent and less stimulatory state, that is, they do not have the properties of mature, stable dendritic cells. If the latter reversion were to take place in vivo during

-4 -

adoptive immunotherapy, the cells could be ineffective as adjuvants. In addition, it is desirable to develop a culture system independent of fetal calf serum.

Engleman et al. International Patent Application WO 95/34638 refers to a method of activating an immune response in a human patient by administering to the patient dendritic cells primed with antigen. Engleman et al. however refer to using dendritic cells isolated from the individual which exist in small numbers. Accordingly, it would be desirable to be able to obtain dendritic cells from a large population of cells present in an easily accessible tissue such as blood.

Two antigens, have recently been reported which, in addition to other antigens or phenotypic characteristics, distinguish mature dendritic cells from other types of white blood cells. Zhou and Tedder (Zhou and Tedder, 1995) reported that CD83, a member of the immunoglobulin superfamily that was cloned from an EBV induced B cell library, is expressed on dendritic cells in blood cells that were cultured for 2 days. This culture period is sufficient to allow a small subset of immature dendritic cells to mature (O'Doherty et al. 1993). CD83 has also been detected on some presumptive dendritic cells in the T cell areas of lymphoid organs, and on some B cells in germinal centers (Zhou et al. 1992). Langhoff and coworkers found that p55, an actin bundling protein, also marks the dendritic cells that are found in 2 day cultures of human blood (Mosialos et al. 1995). p55 is an intracellular protein that was discovered as an EBV induced host cell product. It is expressed by interdigitating cells and at high levels in the brain (Mosialos et al. 1995).

SUMMARY OF THE INVENTION

This invention provides methods of preparing large numbers of stable, mature dendritic cells. The stable mature dendritic cells provided by this invention retain characteristics of a mature phenotype, including expression of dendritic cell markers p55 and CD83 and high levels of accessory molecules like CD86 and CD40, even when removed from contact with cytokines which promote the maturation of pluripotential peripheral blood mononuclear cells (PBMCs) to non-stable immature dendritic cells. As used herein, "stable" refers to retention of a mature dendritic cell phenotype for three days when mature dendritic cells are cultured in the absence of cytokines. Dendritic cells are considered "mature" according to this invention when they have an increased expression of one or more phenotypic markers, such as for example non-adherence to plastic and CD86 or other antigen markers, associated with the accessory function of dendritic cells following contact with a dendritic cell maturation factor.

In one embodiment of the invention, mature dendritic cells are produced in vivo or in vitro from immature dendritic cells derived from PBMC pluripotential cells having the potential of expressing either macrophage or dendritic cell characteristics. The method, according to the invention, comprises contacting the immature dendritic cells with a dendritic cell maturation factor. The dendritic cell maturation factor, may actually be one or more different substances, and may be provided by substances including, but not limited to PBMC conditioned media, maturation factors purified from the conditioned medium, and SACS (fixed Staphylococcus aureus Cowan 1 strain (Pansorbin)). Culture of the pluripotential cells with cytokines such as for example, a combination of GM-CSF and IL-4 or IL-13 stimulates the differentiation of the pluripotential cells to immature dendritic cells.

-6 -

The method provided by this invention is particularly useful with human cells and provides sufficient numbers of mature dendritic cells to be useful for priming with antigen in vitro and activating an individual's T cells by administering the primed dendritic cells to the individual, or activation of T cells in vitro which are then administered to the individual. Accordingly, another aspect of this invention is a method of activating an individual's T cells comprising:

a) obtaining a population of pluripotential cells having the potential of expressing either macrophage or dendritic cell characteristics from an individual and placing them in culture;

b) contacting the pluripotential cells of step a with at least one cytokine to produce immature dendritic cells;

c) contacting the immature dendritic cells with a dendritic cell maturation factor for a time sufficient to cause said immature dendritic cells to stably express dendritic cell characteristics;

d) contacting the dendritic cells obtained from step c with an antigen to produce primed dendritic cells;

e) exposing T cells to the primed dendritic cells of step d to activate said T cells. It is to be understood that one or more of the above steps may be combined so as to occur concurrently, such as for example, steps c and d.

Activation of T cell responses against a wide variety of antigens is facilitated by this invention. Such antigens include, but are not limited to viral, bacterial, tumor and self antigens such as antigen receptors, e.g. T cell receptors. Accordingly, the dendritic cells prepared according to this invention are useful for the prevention



- 7 -

and treatment of various diseases including infectious disease, cancer and autoimmune disease.

The methods of promoting maturation of dendritic cells to mature dendritic cells using cytokines and a dendritic cell maturation factor, and the subsequent activation of T cells, may also be applied to individuals in vivo. According to this embodiment, an individual may be administered cytokines, e.g. GM-CSF and IL-4, or G-CSF to stimulate in vivo the production of immature dendritic cells. A dendritic cell maturation factor, such as conditioned medium, or a substance which stimulates the release of a dendritic cell maturation factor from PBMCs, may also be administered to an individual to stimulate in vivo maturation of immature dendritic cells, the population of which may optionally have been increased through prior administration of cytokines.

This invention also encompasses the mature dendritic cells prepared according to the method of the invention and pharmaceutical compositions comprising the mature dendritic cells and a physiological carrier. Such carriers may include the media described below.

Another aspect of this invention is a culture medium useful for causing the maturation of immature dendritic cells to phenotypically stable, mature dendritic cells. The culture medium of the invention comprises a mixture of salts, carbohydrates and vitamins at physiologic concentrations typical of commercially available culture medium; as well as about 1-5%, and more preferably 1% human serum or plasma; GM-CSF and one or more cytokines which promote maturation of the PBMC phenpotential cells to immature dendritic cells and in amounts together sufficient to promote maturation of pluripotential PBMC to immature dendritic cells; and a sufficient concentration of a dendritic cell maturation factor to cause the maturation of immature dendritic cells to stable mature

-8 -

dendritic cells. This method can also be used with serum-free medium such as for example, XVIVO-20 or AIM-V.

Another aspect of this invention is an assay to detect a dendritic cell maturation factor. Such an assay is useful for the identification and production of the factor present in PBMC conditioned medium which causes the maturation of immature dendritic cells to the mature dendritic cells provided by this invention. This assay comprises contacting a test sample with a culture of unstable, immature dendritic cells derived from a population of pluripotential cells having the potential of expressing either macrophage or dendritic cell characteristics and which express characteristics of immature dendritic cells when cultured in a medium containing at least one cytokine. The presence of a dendritic cell maturation factor in the test substance is then determined by detecting the maturation of the immature dendritic cells in response to the test substance.

It is an object of this invention to provide methods and compositions useful for activating an individual's T cells against specific antigens. The activation of an individual's T cell is useful for the prevention or treatment of disease, for example killer cells to treat or vaccinate against cancer or infection.

It is another of this invention to provide methods useful for causing the stable maturation of immature dendritic cells to mature dendritic cells suitable for administering to an individual for the purpose of activating their immune response, and in particular, T cells.

Another object of this invention is to provide mature dendritic cells cultured in vitro from non-proliferating pluripotential cells.

-9 -

Another object of this invention is to provide culture mediums which promote the maturation of immature dendritic cells to stable mature dendritic cells.

Yet another object of this invention is to provide an assay useful for detecting the presence in a test sample of a dendritic cell maturation factor.

Another object of this invention to provide methods for stimulating the maturation of dendritic cells in vivo.

#### BRIEF DESCRIPTION OF THE FIGURES

10 FIGURE 1: Conditioned medium (CM) is required to ensure the maturation of dendritic cells from progenitor cells. ER- cells were cultured for 7 days in RPMI medium containing 1% plasma in the presence of GM-CSF and IL-4. Cells were then transferred to fresh plates and cultured 15 for 4 days in the presence [open squares] or absence [closed circles] of CM. Control cultures were ER- cells cultured in Teflon beakers for 11 days [open circles]. T cell stimulatory function [primary allogeneic MLR] of the various APC populations is shown.

20 FIGURE 2: Cytofluorographic analysis of dendritic cells grown in the presence of GM-CSF/IL-4 and CM. Dead cells and contaminating lymphocytes were excluded by light scatter properties. Dot plots of the remaining cells are shown. Isotype controls are shown in the left panel.

25 FIGURE 3: Morphology and phenotype of blood derived dendritic cells and macrophages. Dendritic cells [left panel] were generated from normal blood ER- cells in the presence of GM-CSF/IL-4 and CM, cytopspun onto glass slides and stained with a panel of mAbs [see Methods]. Macrophages were syngeneic ER- cells, enriched by plastic adherence and cultured for two weeks in Teflon beakers. CD8 serves as the isotype control antibody. Dendritic 30 35

-10 -

cells are distinguishable from macrophages by their dendritic shapes, lack of CD14 expression, high p55 expression and perinuclear pattern of CD68 expression. Black arrows in the right panel point to presumptive dendritic cells contaminating the macrophage populations.

**FIGURE 4:** Attempts to replace conditioned medium with cytokines. ER- cells were cultured in GM-CSF/IL-4 for 7 days and then transferred to fresh tissue culture plates. The cells were supplemented with 50% CM [black circles] in the absence or presence of polyclonal neutralizing antibodies to TNF $\alpha$ , IL-1 $\beta$  and IL-6 [1-5  $\mu$ g/ml]. Cocktail refers to the combination of the various antibodies. Some cultures received TNF alpha [20 ng/ml] or IL-1 beta [10ng/ml] instead of CM. After 4 days [day 11 of culture], the cells were evaluated for T cell stimulatory activity in an allo MLR. Results are shown as  $^3\text{H}$ -TdR incorporation [cpm  $\times 10^3$ ] and are averages of triplicates.

**FIGURE 5:** Dendritic cell progenitors are enriched in the CD14 hi population. ER- cells from two different sources were sorted into CD14 hi and lo populations as described in Methods. Unsorted and various sorted fractions were analyzed for T cell stimulatory activity in an MLR. In [A], the CD14 hi cells were irradiated with 3,000 rads, 137 Cs source. In [B], the CD14 hi cells were irradiated with 1,500 rads. Results are shown as  $^3\text{H}$ TdR incorporation [cpm  $\times 10^3$ ] and are averages of triplicates.

**FIGURE 6:** Morphological stability of mature dendritic cells. Lymphocyte-depleted PBMC were cultured for 10 days in FCS-containing RPMI medium in the presence of GM-CSF and IL-4. Phase contrast micrographs were taken to illustrate the morphology of the cells. [A] No CM was added during the 10 day culture. Cells were washed on day

-11 -

10 and recultured without cytokines for 3 additional days. They re-adhere firmly and have the appearance of macrophages. [B] CM was added from day 7-10. The cells acquire many lamellipodia or veils and are nonadherent. [C,D] CM was added from day 7-10, the cells were washed and returned to culture for 3 days without CM or cytokines. The morphology of mature dendritic cells persists both in RPMI/10% FCS [C] and X-VIVO 20/serum-free [D] media. A, x275; B-D, x550.

**FIGURE 7:** Comparative cytofluorographic analysis of mature vs. immature dendritic cells. Lymphocyte-depleted PBMC were cultured for 7 days with GM-CSF and IL-4 followed by a maturation period from day 8 to day 10 in the presence of CM. The resulting fully mature dendritic cells were compared to corresponding immature dendritic cells that had not been treated with CM. In all cases mature dendritic cells express CD83 and enhanced levels of CD86, and they lose CD115 expression. [A,A'] Dendritic cells cultured from day 7 to day 10 without [A] or with [A'] CM in RPMI/FCS are shown. Fluorescence on the x-axis represents HLA-DR expression; y-axis shows the indicated antibody staining. Dot plots are from cells gated on the basis of their light scatter properties [FSC/SSC]. [B,B'] Cells were analyzed on day 7[B] and on day 10, i.e., after maturation in the presence of CM [B']. Culture medium was RPMI/1% human plasma; lymphocytes were depleted with immunomagnetic beads. Histograms are from cells gated on the basis of their light scatter properties [FCS/SSC]. [C,C' - left] Like B,B', but lymphocytes were initially depleted by E-rosetting. [C,C' - middle] like B,B', but AIM-V medium/1% human plasma was used. [C,C' - right] like A,A', co-2b, isotype-matched control IgG2b [for CD83, 86, 14]; co-rat, control for rat IgG1 [for CD115]; FSC/SSC, forward scatter/side scatter. All markers are

-12 -

set according to staining with isotype-matched control antibodies.

**FIGURE 8:** Fixed staphylococci [SACS] are required to ensure the maturation of dendritic cells from progenitor cells. Lymphocyte-depleted PBMC were cultured for 7 days in RPMI medium supplemented with FCS and GM-CSF/IL-4. Cells were then transferred to fresh plates and cultured for 3 more days in the presence [closed squares] or absence [open squares] of SACS. T cell stimulatory function in the primary allogeneic MLR of the dendritic cells matured in SACS containing medium is markedly enhanced. Neither IL-1 $\beta$  [in A] nor IL-15 [in B] that were given instead of SACS could fully reconstitute the effect.

**FIGURE 9:** Morphology and motility of mature dendritic cells grown in the presence of GM-CSF/IL-4 and CM. Dendritic cell progenitors were primed for 7 days in the presence of GM-CSF and IL-4 followed by a 3 day maturation period in the presence of CM. Culture medium was RPMI supplemented with 1% autologous human plasma. Note the numerous thin cytoplasmic processes ("veils"). Photographs of the same field taken at intervals of 15 seconds show that the veils move. Magnification: x550.

**FIGURE 10:** Phagocytosis of latex beads. Dendritic cells were grown in the presence of GM-CSF and IL-4 until day 8. GCS-containing RPMI medium was used. Cells in the upper panel were exposed to latex beads for 24 hours in the absence of CM. Latex is taken up into the cells (arrows). In the lower panel cells were first allowed to mature in the presence of CM from day 7 to day 10 and were then exposed to latex. Little or no latex is phagocytosed into mature, "veiled" cells. Only a contaminating adherent cell has accumulated beads. x300.

-13 -

FIGURE 11: Immunostimulatory functions of mature dendritic cells. Dendritic cells were cultured in the presence of GM-CSF and IL-4 until day 7 and further on until day 10 with [closed symbols] or without [open symbols] CM. RPMI was supplemented with FCS in A and D-F, and with 1% autologous human plasma in B and C. [A] Note pronounced enhancement of T cell stimulatory capacity in the MLR following exposure to CM. Two independent experiments are shown. [B] Fully mature dendritic cells grown in RPMI supplemented with 10% FCS or 1% autologous human plasma were compared to each other. [C] Fully mature dendritic cells efficiently stimulate naive cord blood T cells. [D] Tetanus toxoid protein [1  $\mu$ g/ml] is presented to a peptide-specific T cell clone. CM-induced maturation leads to a decrease in antigen processing capacity. Triangles in D indicate the [lack of] response of clone cells to APC in the absence of antigen. Values are means of triplicate wells and are expressed as cpm  $\times 10^3$ . [E,F] Peptide-pulsed mature dendritic cells elicit T cell lines that lyse autologous target cells in a peptide dose-dependent [E] and MHC-restricted [F] manner. anti-cl. II, anti-cl. I, anti-MHC class II and I mAb's W6/32 and L243, respectively.

FIGURE 12: Dendritic cell progenitors reside in a CD34-negative population. PBMC of a cancer patient who had been treated with G-CSF were passed through an anti-CD34 immunoaffinity column. The original unseparated population [solid lines] and the CD34-depleted population [dashed lines] were cultured in RPMI\FCS with GM-CSF and IL-4 until d7 and further on from d7 to d10 in the presence [closed symbols] or absence [open symbols] of CM. Both populations gave rise to dendritic cells that matured upon exposure to CM.

-14 -

DETAILED DESCRIPTION OF THE INVENTION

This invention relates to methods and compositions useful for promoting the maturation of immature dendritic cells to a mature and stable phenotype as well as to the stable mature dendritic cells. The mature dendritic cells are useful as antigen presenting cells (APCs) which activate other immune cells including antigen specific helper and killer T cells. Thus, this invention also provides methods and compositions useful for the treatment of disease.

The method of producing mature dendritic cells according to this invention comprises contacting immature dendritic cells with a dendritic cell maturation factor. We have found such a dendritic cell maturation factor in conditioned medium obtained from cultures of peripheral blood mononuclear cells (PBMCs). Immature dendritic cells may be derived from PBMCs by culturing PBMCs with cytokines, such as, for example, a combination of GM-CSF and IL-4, which promote their differentiation to immature dendritic cells. Surprisingly, we have determined that unless they are exposed to a dendritic cell maturation factor, removal of the cytokines from contact with the cells causes the cells to revert back to a pluripotential cell having characteristics similar to macrophages.

The response of immature dendritic cells to contact with the maturation factor is an increased expression of CD83 and p55 dendritic cell markers; strong expression of antigen presenting MHC class I and II products; and expression of several accessory (adhesion and co-stimulatory) molecules including CD40, CD54, CD58, CD80 and CD86. A decrease in expression of CD115 as well as CD14, CD68 and CD32, which are markers associated with immature dendritic cells, is also observed as a result of carrying out the method of this invention. In addition, this phenotype remains stable for up to three days even



-15 -

after removal of cytokines used to promote cell maturation.

#### Sources of Pluripotential Cells

5 The pluripotential cells, from which the immature dendritic cells for use in this invention are derived, are present in blood as PBMCs. Although most easily obtainable from blood, the pluripotential cells may also be obtained from any tissue in which they reside, including bone marrow and spleen tissue. These pluripotential cells typically express CD14, CD32, CD68 and CD115 monocyte markers with little or no expression of CD83, p55 or accessory molecules such as CD40 and CD86. When cultured in the presence of cytokines such as a combination of GM-CSF and IL-4 or IL-13 as described below, the pluripotential cells give rise to the immature dendritic cells.

15 Methods of obtaining PBMCs from blood, such as differential sedimentation through an appropriate medium, e.g. Ficoll-Hypaque [Pharmacia Biotech, Uppsala, Sweden], are well known and suitable for use in this invention. In a preferred embodiment of the invention, the pluripotential cells are obtained by depleting populations of PBMCs of platelets, and T and B lymphocytes. Various methods may be used to accomplish the depletion of the non-pluripotential cells. According to one method, immunomagnetic beads labelled with antibodies specific for T or B lymphocytes, either directly or indirectly may be used to remove the T and B cells from the PBMC population. T cells may also be depleted from the PBMC population by rosetting with neuraminidase treated red blood cells as described by O'Dherty (1993), which is incorporated herein by reference.

30 To produce 3 million mature dendritic cells, it is necessary to process about 40 mls of blood. 4 to 8 x 10<sup>7</sup>

35

-16 -

pluripotential PBMC give rise to approximately 3 million mature dendritic cells.

Culture of Pluripotential PBMCs To Produce Immature Dendritic Cells

5       The immature dendritic cells for use in this invention are post-mitotic but not yet terminally differentiated to either a macrophage or dendritic cell phenotype. Cultures of immature dendritic cells may be obtained by culturing the pluripotential cells in the presence of cytokines which promote their differentiation. A combination of GM-CSF and IL-4 at a concentration of each at between about 200 to about 2000 U/ml, more preferably between about 500 and 1000 U/ml, and most preferably about 800 U/ml (GM-CSF) and 1000 U/ml (IL-4) produces significant quantities of the immature dendritic cells. A combination of GM-CSF (10 ng/ml) and IL-4 (10-20 ng/ml) has been found to be useful with this invention. It may also be desirable to vary the concentration of cytokines at different stages of the culture such that freshly cultured cells are cultured in the presence of higher concentrations of IL-4 (1000 U/ml) than established cultures (500 U/ml IL-4 after 2 days in culture). Other cytokines such as IL-13 may be found to substitute for IL-4.

25       Methods for obtaining these immature dendritic cells from adherent blood mononuclear fractions are described in Romani et al. (1994); and Sallusto and Lanzavecchia, 1994) both of which are incorporated herein by reference. Briefly, lymphocyte depleted PBMCs are plated in tissue culture plates at a density of about 1 million cells/cm<sup>2</sup> in complete culture medium containing cytokines such as GM-CSF and IL-4 at concentrations of each at between about 800 to 1000 U/ml and IL-4 is present at about 1000 U/ml.

35

-17 -

Various mediums are suitable to initially culture PBMCs. Typical media for use in this invention comprise physiologic inorganic salts, carbohydrates including sugars, amino acid, vitamins and other components known to those in the art. Preferred media formulations include, for example, RPMI 1640, X-VIVO 20, AIM-V, HybriCare and Iscove's. More preferably the medium is RPMI 1640 or X-VIVO 20 or AIM-V. Most preferably, the medium is RPMI 1640.

It is desirable to avoid the use of fetal calf serum (FCS) in cultures for human use. The presence of bovine proteins may cause unwanted sensitization of the recipient against bovine proteins and it is therefore desirable to substitute FCS with human serum or plasma in an amount sufficient to maintain cell viability. The use of autologous, non-heat-inactivated human plasma at a concentration of about 1% is preferred. Using RPMI 1640 supplemented with 1% autologous human plasma, a final yield of between about  $0.8$  to  $3.3 \times 10^6$  CD83+ mature dendritic cells is obtainable from 40 ml of blood.

Another source of immature dendritic cells are cultures of proliferating dendritic cell precursors prepared according to the method described in Steinman et al. International application PCT/US93/03141, which is incorporated herein by reference. Since the dendritic cells prepared from the CD34+ proliferating precursors mature to dendritic cells expressing mature characteristics it is likely that they also pass through a development stage where they are pluripotential. By treating these cultures with a dendritic cell maturation factor as described below, the efficiency of obtaining mature dendritic cells from proliferating precursors may also be improved.

### Maturation of Immature Dendritic Cells

The mature dendritic cells are prepared according to this invention by contacting the immature dendritic cells with a dendritic cell maturation factor. As referred to herein, the dendritic cell maturation factor may actually be one or more specific substances which act in concert to cause the maturation of the immature dendritic cells. Such a factor has been determined to be present in PBMC conditioned medium, preferably monocyte conditioned medium. Another means of obtaining dendritic cell maturation factors are to treat PMBCs with substances which stimulate the release of such factors. Substances such as immunoglobulin or SACS (also referred to as Pansorbin) are useful for causing monocytes to release cytokines.

After the cells have been cultured for a sufficient time to express some characteristics of dendritic cells, preferably about 6 to 10 days, PBMC conditioned media is added to the culture medium at a final concentration of between about 10 to 50%. More preferably between about 25 and 50%. Maturation of the immature dendritic cells to the mature phenotype may occur without the presence in the medium of the cytokines used to stimulate maturation of the pluripotential cells to immature dendritic cells. Typically, the immature dendritic cells are obtained after about 6 to 7 days in culture with the cytokines. Preferably the immature dendritic cells are subcultured prior to addition of the dendritic cell maturation factor, e.g., conditioned medium.

In a preferred embodiment, the immature dendritic cells which are non-adherent cells are harvested and subcultured in the presence of the dendritic cell maturation factor present in the conditioned medium, and optionally with cytokines such as GM-CSF and IL-4. The mature dendritic cells are typically obtained within

-19 -

approximately three days following addition of the conditioned medium.

Variations in the culture protocol which are also considered within the scope of this invention may include, but not be limited to, changes in cytokines used to promote differentiation of PBMCs to immature dendritic cells, concentrations of cytokines and timing of the addition of the dendritic cell maturation factor present in the conditioned medium.

#### Characterization of Conditioned Medium

The conditioned medium for use in this invention may be made by culturing PBMCs, preferably monocytes in basic growth medium. Preferably the cells for producing conditioned medium are cultured in the presence of a stimulus which stimulates the release of factors from the donor cells. An example of such a preferred factor is gamma globulin when bound to the culture substrate. Methods for producing PBMC (monocyte) conditioned medium have previously been described in O'Doherty et al. (1993 and 1994) which are incorporated herein by reference. PBMC or preferably T cell depleted, or more preferably T and B cell depleted are cultured (about  $5 \times 10^7$  cells/100 mm plate) for about 24 hours in medium (6-8 mls) containing 1% of plasma, preferably autologous plasma. Where conditioned medium is to be used for stimulating maturation of dendritic cells in vivo, the growth medium should be selected to be compatible with use in humans, e.g. X-VIVO or AIM-V and preferably is used without serum or plasma. For use in humans the conditioned medium is concentrated using standard procedures and adjusted to be isotonic so as to be suitable for administration in vivo. Using this method, equivalent concentrations of dendritic cell maturation factor can be achieved in vivo, as in vitro.

-20 -

In addition to gamma globulin as a stimulant of cytokine release from the culture PBMCs, other cytokine stimulators such as SACS (fixed Staphylococcus Aureus Cowan 1 strain, 2,01 mg/ml Ig-binding capacity: Pansorbin cells, Cat. No. 507861) may also be used. Because SACS is bacterially derived its use is less desirable than Ig.

In addition to stimulating the production of cytokines for producing conditioned medium, SACS may also be substituted for conditioned medium as a dendritic cell maturation factor. When used to promote maturation of the immature dendritic cells, SACS is added to cultures at a concentration sufficient to cause their maturation, preferably at about 1:10000 dilution.

This invention provides evidence for a new pathway for dendritic cell development. Prior studies have indicated that dendritic cells arise either from CD34+ proliferating progenitors (Inaba et al. 1992; Caux et al. 1992), or from nonproliferating "null" cells that lack monocyte and lymphocyte markers [CD3, CD14, CD16, CD19] (O'Doherty et al. 1993). Both of these progenitor populations are infrequent in human blood representing <0.1% and <1% of blood mononuclear cells, respectively. Sallusto and Lanzavecchia [personal communication] reason that monocytes can give rise to dendritic cells in the presence of FCS, but we find that these cells are not fully mature in several respects. In contrast, large numbers of mature dendritic cells can develop, and in human plasma or serum, when CM is added to the GM-CSF and IL-4 priming step. The precursors are radioresistant and primarily in the standard "monocyte" fraction [CD14 positive and plastic adherent]. It is possible that most monocytes are bipotential and can develop into either macrophages or dendritic cells depending upon the cytokines that are applied. Alternatively, only some of the "monocyte" fraction may have the potential to develop

-21 -

into dendritic cells. At this time, our data are consistent with the latter possibility since the total yield of dendritic cells from radioresistant precursors represents about 5.5-16% of the starting number of monocytes.

Physiologic counterparts for the maturation over several days of dendritic cells from nonproliferating blood progenitors can be proposed in at least three settings. First sizable numbers of dendritic cells traffic in afferent lymphatics, and these cells derive from precursors that have been proliferating 3-7 days prior to detection in lymph (Mason et al. 1981; Pugh et al. 1983; Fossum, 1989a; Fossum, 1989b). These afferent lymph dendritic cells may derive from blood progenitor cells that undergo a priming phase. Second, dendritic cells are rapidly induced to migrate into tissues [lung, gut] in response to pro-inflammatory stimuli e.g. LPS or infection (Watson et al. 1990; McWilliam et al. 1994). There they may undergo further maturation in response to cytokines that are produced locally by tissue macrophages.

Third, significant numbers of mature dendritic cells accumulate in rheumatoid synovial exudates during acute flares of arthritis (Zvaifler et al. 1985; Bhardwaj et al. 1988; Helfgott et al. 1988; Thomas et al. 1994). These effusions are rich in GM-CSF, IL-4, IL-6 and TNF alpha, amongst other cytokines [reviewed in (Klareskog et al. 1995)]. In addition, the presence of substantial numbers of macrophages and immune complexes may provide a natural "conditioned medium" for the maturation and differentiation of dendritic cells from blood progenitors that enter inflamed joints.

Based on the data reported in Examples 1 and 2, dendritic cells cultured from peripheral blood precursors in GM-CSF and IL-4 for 6-7 days can be considered as immature dendritic cells. Although they have already

-22 -

acquired relatively high levels of MHC and adhesion/costimulator molecules as well as the capacity to stimulate resting T cells they still lack markers for terminal dendritic cell differentiation such as CD83. They are however well equipped with the necessary prerequisites for the processing of native protein antigens. They can phagocytose particulate matter and they can efficiently process native tetanus toxoid protein into immunogenic MHC-peptide complexes recognized by a tetanus peptide-specific T clone. In addition, Sallusto et al. (1995) described active uptake of soluble macromolecules by macropinocytosis, and we find uptake of latex particles as well. Immature dendritic cells may therefore be useful in immunization protocols that use native proteins or particles as antigens. One must keep in mind, however, that the immature dendritic cells generated with GM-CSF and IL-4, but without CM, are not stable. Upon withdrawal of cytokines they re-adhere and appear to revert back to monocytes. Such cells would therefore not be as useful for therapeutic approaches. It will be preferred to induce maturation of antigen-pulsed dendritic cells by the method described here, i.e., exposure to monocyte-conditioned media.

Dendritic cells grown in GM-CSF and IL-4 for 7 days and exposed to macrophage-conditioned media for another three days develop into mature dendritic cells. These cells have down-regulated their antigen uptake mechanisms and their processing capacity. CM- treated dendritic cells correspond morphologically, phenotypically and functionally to well defined populations of mature dendritic cells, e.g., cultured epidermal Langerhans cells (Romani et al. 1989b); Tenunissen et al. 1990), cutaneous dendritic cells obtained by emigration from skin explants (Pope et al. 1995; Lenz et al. 1993), or blood dendritic cells obtained by classical methods involving 36 hours of



-23 -

culture (Young and Steinman, 1988). They are specialized in the sensitization of naive, resting T cells. We show here that they can efficiently induce proliferation in allogeneic umbilical cord T cells which may be considered virgin T cells. We also demonstrate that these mature dendritic cells can be loaded with an antigenic peptide and induce antigen-specific cytotoxic T lymphocytes from populations of autologous PBMC or CD8+ T cells. In addition, mature dendritic cells in small numbers are able to elicit rapid and virus-specific CTL responses from quiescent autologous T cells. Mature dendritic cells will be helpful in immunization protocols that employ antigenic peptides preferably to proteins. This applies especially to the field of tumor immunotherapy. Tumor-specific peptides are being discovered at a fast rate (Boon et al. 1994).

Based on our data, the optimal method to generate mature dendritic cells for clinical purposes would be to deplete PBMC of lymphocytes with immunomagnetic beads, culture them for 6 to 7 days in RPMI medium supplemented with 1% autologous human plasma and GM-CSF/IL-4, and induce them to mature within another three days of culture by the addition of conditioned medium that is produced by PBMC adhering to Ig-coated Petri dishes. All these reagents are already approved [Anti-mouse Ig Dynabeads and mouse anti-T and B cell mAb's/Baxter; GM-CSF/Sandoz; IL-4/Schering-Plough; Ig/Biochemie-Sandoz] or have been used in clinical studies [RPMI medium, Lymphoprep].

Our method to generate dendritic cells from progenitor cells in autologous plasma will be useful in several respects. First, the method is simple, reproducible and generates  $1-3 \times 10^6$  dendritic cells from relatively small, 40-50 ml blood samples. Second, critical features of dendritic cells such as pathways of antigen presentation and T cell signalling can be studied

-24 -

in the absence of foreign FCS derived proteins. Third, one has ready access to cells that can be used as adjuvants to enhance protective immune responses in vivo. We find that dendritic cells generated by the method described here and pulsed with either influenza virus [live or heat inactivated] or immunodominant peptides [Table 6] induce virus-specific CD8+ killer cell responses in vitro.

#### Pulsing Dendritic Cells With Antigen

The mature dendritic cells prepared according to this invention are useful for activating T cells against specific antigens. Several antigens and methods for priming dendritic cells have been described and may be adapted for use in this invention. See, for example Engleman et al. International patent application PCT/US95/07461; Hsu et al., Nature Med. 2:52-58 (1996); and Steinman et al. International application PCT/US93/03141 which are all incorporated herein by reference. Antigens associated with fungal, bacterial, viral, tumor, or autoimmune (i.e., self antigens) diseases are useful for priming dendritic cells to activate T cells which aid in treating or preventing disease.

Where it is desirable for cells to take up antigen by phagocytosis, it is preferable to add antigen to the cultures of immature dendritic cells prior to addition of the dendritic cell maturation factor. Phagocytosis may be desirable when particulate antigens, or immune complexes are used. In most cases it is sufficient to expose antigen to the dendritic cells after they have attained the mature phenotype or while they are exposed to the dendritic cell maturation factor to attain the mature phenotype. This method is preferred when soluble peptide antigens are used.

-25 -

For the purpose of priming cells typically approximately  $1$  to  $5 \times 10^6$  cells are exposed to antigen at a concentration of between about  $10$  pM to about  $10$   $\mu$ M, inclusive. More preferably about  $1$   $\mu$ M antigen to about 3 million cells is used. The dendritic cells are cultured in the presence of the antigen for a sufficient time to allow for uptake and presentation. Typically uptake and processing can occur within 24 hours but longer (up to and including 4 days) or shorter (about 1-2 hours) periods may also be used.

For activating T cells in an individual between about  $2 \times 10^5$  and  $2 \times 10^9$  more preferably between about 1 million and 10 million mature dendritic cells should be administered to the individual. The dendritic cells should be administered in a physiologically compatible carrier which is nontoxic to the cells and the individual. Such a carrier may be the cell growth medium described above. The mature dendritic cells prepared according to this invention are particularly potent at activating T cells. For example, using prior methods of obtaining dendritic cells the ratio of dendritic cells to T cells necessary for strong T cell activation is about 1 dendritic cell to 30 T cells whereas the ratio according to this invention is about 1 to 1000. Thus, fewer dendritic cells are required. For activating T cells in vitro the ratio of dendritic cells to T cells is between about 1:10 and 1:1000. More preferably between 1:30 and 1:150. Between approximately  $10^8$  and  $10^9$  activated T cells are administered back to the individual to produce a response against an antigen.

Dendritic cells may be administered to an individual using standard methods including intravenous, intraperitoneal, subcutaneously, intradermally or intramuscularly. The homing ability of the dendritic

35

-26 -

cells facilitates their ability to find T cells and cause their activation.

The methods of this invention are particularly well suited for use against tumors from which tumor specific antigens are obtainable or which express a mutated protein. Preferrably, the antigen is a molecule required by the tumor cells to be malignant, for example altered ras. Non-limiting examples of tumors for which tumor specific antigens have been identified include melanoma, B cell lymphoma, uterine or cervical cancer. An example of a melanoma antigen which could be considered for use to prime dendritic cells is the human melanoma specific antigen gp75 antigen (Vijayasardhi S. et al., J. Exp. Med., 171:1375-1380 (1990). An example of an antigen useful for targeting cells against cervical cancer is papilloma virus. Tumor specific idiotypic protein derived from B cell lymphomas has been reported to prime dendritic cells which activate T cells against the tumor cells. Hsu et al. (1966), supra. In addition to tumor antigens, viruses may also be used to prime the dendritic cells. Viruses may also be used to cause dendritic cells to present viral antigens, or they may be engineered to express non-viral proteins which are then processed and presented by the dendritic cells. Non-limiting examples of viruses which may be used to prime dendritic cells include, but are not limited to, influenza, HIV, CMV, EBV, human papilloma virus, adenovirus, HBV, HCV and vaccinia. Isolated viral proteins, inactivated or attenuated virus may also be used. Non-limiting examples of bacterial or protozoan antigens include tetanus toxoid, BCG, malaria antigens and leishmania antigens. Specific antigens associated with autoimmune disease such as immune receptors may also be used to prime dendritic cells to be used to activate T cells against such antigens.

35

-27 -

o  
Method of Assaying for a Dendritic Cell Maturation Factor

This invention provides an assay useful for identifying the dendritic cell maturation factor present in a test substance such as monocyte conditioned medium. This assay is based on the ability of immature dendritic cells to mature and express a stable phenotype after having been contacted with a dendritic cell maturation factor. Thus, the assay comprises contacting a culture of immature dendritic cells with a test substance and detecting the maturation of the immature dendritic cells in response to the presence of the test substance. The immature dendritic cells for use in this assay are derived from a population of pluripotential cells, such as human PBMCs having the potential of expressing either macrophage or dendritic cell characteristics and which express characteristics of immature dendritic cells when cultured in a medium containing cytokines (e.g., GM-CSF and IL-4). Maturation associated with the presence of a dendritic cell maturation factor may be confirmed, for example, by detecting either one or more of 1) an increase expression of one or more of p55, CD83, CD40 or CD86 antigens; 2) loss of CD115, CD14, CD32 or CD68 antigen; or 3) reversion to a macrophage phenotype characterized by increased adhesion and loss of veils following the removal of cytokines which promote maturation of PBMCs to the immature dendritic cells. The assay described above may be used in conjunction with biochemical techniques to identify the dendritic maturation factor present in monocyte CM. In addition, it provides a useful bioassay to confirm the activity of such a factor once it is identified or purified in connection with its manufacture.

-28 -

EXAMPLESEXAMPLE 1MethodsCulture medium

RPMI 1640 supplemented with 20  $\mu\text{g/ml}$  of gentamicin, 10 mM HEPES, and either 10% heat-inactivated FCS [Gemini Bioproducts, Calabasas, Ca] or 1% and 10% autologous plasma [heparinized].

Cytokines

We were generously supplied with rhGM-CSF [ $1 \times 10^5$  U/ $\mu\text{g}$ ] by Kirin Brewery Co., Maebashi, Gunma, Japan, and rhIL-4 [ $0.5 \times 10^5$  U/ $\mu\text{g}$ ] by Immunex Corp. rhIL-12 [ $2 \times 10^7$  U/mg] was provided by the Genetics Institute, Boston, Mass. The recombinant human cytokines IL-15, IL-1, IL-6, TNF alpha were purchased from R and D Corp [Minneapolis, Mn].

Generation of dendritic cells from human blood

Peripheral blood was obtained from normal donors in heparinized syringes, and PBMC isolated by sedimentation in Ficoll-Hypaque [Pharmacia Biotech, Uppsala, Sweden]. T cell-enriched [ER+] and T cell-depleted [ER-] populations were prepared by rosetting with neuraminidase treated-sheep red blood cells as described (Carr and Kim, 1993). Different starting populations of PBMC were plated in 3 ml volumes in 6 well tissue culture dishes [Falcon] in complete medium. They consisted of [a] plastic adherent and nonadherent PBMC [obtained after 60 min adherence of  $6 \times 10^6$  PBMC/well]; [b] ER- cells; [c] ER- cells sorted on a FACStar<sup>™</sup> into CD14 high or low cells. Groups [b] - [c] were plated at  $2 \times 10^6$ /well. GM-CSF and IL-4 were added at final concentrations of 1000 U/ml on the initial day of culture. Cytokines were replenished

-29 -

every other day [days 2, 4 and 6] by removing 0.3 ml of the medium and adding back 0.5 ml fresh medium with cytokines. On day 7, non-adherent cells were collected by moderately vigorous aspiration and analyzed immediately, maintained in culture or transferred to new 6 well plates. When cultured beyond 7 days, the cultures were supplemented with Pansorbin, [1:10,000 dilution of 2 mg IgG/ml, Calbiochem, La Jolla, Ca], or conditioned medium [final concentration 33 or 50% v/v, see below] at day 7.

#### Conditioned medium

Ig coated bacteriologic plates [100 mm, Falcon, Lincoln Park, N.J.] were prepared immediately prior to use by the addition of 4 ml of human gamma-globulin [10 mg/ml, Cappel Labs, Organon Teknika, Westchester, Pa] for 1 min. The residual gamma-globulin was removed and saved for reuse at least 4-6 times. The plates were washed three times with phosphate buffered saline [PD] prior to use. T cell-depleted ER- cells [ $5 \times 10^7$ ] were layered onto the Ig-coated bacteriologic plates for 1 hour in volumes of 6-8 ml. Non-adherent cells were washed off with gentle aspiration. The gamma-globulin adherent cells were incubated in fresh complete medium with 1% autologous plasma at 37°C for 24 h. The medium was collected and frozen at -20°C prior to use. In some cases, conditioned medium was prepared from cells stimulated with pansorbin instead of human Ig.

#### Monoclonal Antibodies [mAbs]

MABs towards the following antigens were used: HLA-DR, CD14, CD3, CD4/CD8, CD5, CD19, CD16, CD56, CD1a, CD54, CD58, CD25, CD45RO, CD11a, CD11b and CD11c, [Becton-Dickinson, Mountainview, CA], CD80 and CD86 [IgG1, FITC conjugate, Pharmingen], CD83 [IgG1, PE conjugate, Coulter Corp, Miami, Florida], CD68 [Dako, Carpinteria, Ca], CD40,

-30 -

CD95/fas [Immunotech, Marseilles, France], p55 [K-2 clone, a gift from Dr. E. Langhoff; Mosialos et al. (1996)], mannose receptor [mAb 3.2PB1, a gift from Dr. A. Lanzavecchia], Lag anti-Birbeck granule associated antigen [a gift from Dr. S. Imamura], Ki 67 [MIB-1, IgG1, Dako]. Secondary antibody was PE-conjugated F(ab')<sub>2</sub> goat anti-mouse IgG [gamma and light chain, Tago, Burlingame, Ca].

#### Polyclonal neutralizing antibodies

Goat anti-human polyclonal antibodies were purchased from R and D Corp, Minneapolis, MN. They included antibodies toward TNF $\alpha$ , IL-1 beta, IL-1 alpha, IL-12 and IL-6. Control antibody was goat IgG [R and D Corp.].

#### Cytofluorography and cell sorting

Cell populations were phenotyped with the panel of Mabs listed above and analyzed on a FACScan. To obtain CD14<sup>+</sup> cells, 2 x 10<sup>7</sup> ER<sup>-</sup> cells were stained with PE-CD14 [Becton-Dickinson, Mountainview, CA] for 30 min on ice, washed extensively and then sorted on a FACStar<sup>Plus</sup>. Both CD14 high and CD14 low cells were evaluated for the ability to develop into dendritic cells.

#### Immunohistochemistry

Cytospins of various cell populations were prepared using a Cytospin 2 centrifuge [Shandon, Inc., Pittsburgh, PA]. Slides were fixed in absolute acetone for 5 min at room temperature and then incubated with mAbs for 45 min. Cytospins were then washed and incubated with 1:200 dilution of biotinylated goat anti-mouse IgG [Boehringer Mannheim Biochemicals, Indianapolis, IN] for 45 min, followed by a horseradish peroxidase [HRP]-biotin-avidin complex [ABC kit; Vector laboratories, Inc., Burlingame, CA] for 30 min. Non-bound HRP was then washed off, and



-31 -

the HRP reaction product was developed with H<sub>2</sub>O<sub>2</sub> and diaminobenzidine tetrahydrochloride [Polysciences, Warrington, PA].

#### 5 Allogeneic MLR

To test for T cell stimulatory function, the APCs were unirradiated or irradiated [Gy 30] and added in graded doses as stimulators for 2 x 10<sup>5</sup> purified, allogeneic T cells in 96 well flat bottomed plates [Costar]. Proliferation was determined on days 4-6 with  
10 the addition of 4 µCi/ml of [<sup>3</sup>H]TdR for 10-16 h to triplicate wells [mean cpm].

#### Induction of influenza virus specific CTL

Dendritic cells prepared from HLA A2.1+ donors were  
15 washed out of medium containing serum and resuspended in RPMI to 0.5 - 1 x 10<sup>7</sup> cells/ml. Live or heat-inactivated influenza virus [PR8, Spafas Inc., Storrs, Ct] was added at a final concentration of 1,000 HAU/ml for 1 hr at 37°  
20 as previously described (Bhardwaj et al. 1994; Bender et al. 1995). Following three washes, 3 x 10<sup>4</sup> dendritic cells were added to purified T cells [1 x 10<sup>6</sup>] in 48 well plates [Costar, Cambridge, Mass.] After 7 days, the T cells were assayed for cytolytic activity on uninfected or  
25 influenza-infected autologous macrophage targets using a conventional <sup>51</sup>Cr release assay (Bhardwaj et al. 1994). Alternatively, targets were HLA A2.1 matched T2 cells [the TAP deficient line, provided by Dr. P. Cresswell] pulsed for 1 hr with 10 nM influenza virus matrix peptide  
30 [GILGFVFTL]. Target cells were labeled with Na<sup>51</sup>CrO<sub>4</sub> as previously described (Bhardwaj et al. 1994).

#### RESULTS

Cells Cultured in the presence of GM-CSF and IL-4  
35 only were not fully mature. We first confirmed that GM-

-32 -

CSF and IL-4 induces the development of large numbers of potent APCs from lymphocyte-depleted PBMC when cultured for 6-7 days in the presence of FCS [ $2.5 \times 10^6$  per 40 ml of blood, Table 1 first row].

TABLE 1

Allogeneic MLR

$^3\text{H}$ -TdR Incorporation, cpm  $\times 10^{-3}$  AT T:APC ratios of:

Treatment	100:1	300:1	900:1	DC Yield/40ml blood	% Enrichment
10% FCS	153 [17]	33 [3]	7	$2.5 \times 10^6$	6%
10% FCS + PS	254 [196]	154 [135]	79	$3.5 \times 10^6$	95%
10% FCS + CM	258 [55]	129 [27]	39	$3.8 \times 10^6$	96%
1% Plasma	175 [3]	68 [1]	8	$0.5 \times 10^6$	25%
1% Plasma + PS	309 [154]	161 [80]	56	$2.0 \times 10^6$	75%
1% Plasma + CM	286 [8]	156 [1]	48	$1.5 \times 10^6$	74%

Table 1: T cell stimulatory activity of dendritic cells derived from progenitors in blood

ER- cells were cultured for 7 days in RPMI medium containing GM-CSF and IL-4 and either 10% FCS or 1% human plasma. The cells were then cultured for 4 more days in the absence or presence of conditioned medium prepared from pansorbin or gamma globulin stimulated blood mononuclear cells [50% final concentration]. T cell stimulatory function was assessed in the primary allogeneic MLR. Results are shown as  $^3\text{H}$ -TdR incorporation [cpm  $\times 10^{-3}$ ] and are averages of triplicates. \* Numbers in parentheses represent cpm measured with syngeneic T cells. The yield is expressed as the total number of dendritic cells obtained from an original donation of 40 ml of blood. The % enrichment is defined as the percentage of cells in the non-adherent fraction that display typical features of dendritic cells: stellate shape, high expression of HLA-DR, the dendritic cell restricted marker CD83 [see Fig. 2].

-33 -

We discovered, however, that dendritic cells primed under these conditions were not fully mature. Upon immunolabeling of cytopspins or cell suspensions [FACS], there was residual expression of the monocyte markers CD14 and CD32 and weak to no expression of the dendritic cell restricted markers CD83 (Zhou et al. 1992; Zhou and Tedder, 1995) and p55 (Mosialos et al. 1996).

Furthermore, when FCS primed cells were recultured without cytokines, MLR stimulating activity decreased considerably and most of the cells attached firmly to plastic [data not shown]. These findings suggested to us that GM-CSF and IL-4 were insufficient triggers to generate mature and stable dendritic cells from non-proliferating blood progenitors.

When 10% human serum or plasma was used in the GM-CSF/IL-4 priming culture in the place of FCS, most of the PBMC remained adherent and few free dendritic cells developed by 7 days. To reduce adherence, medium containing 1% plasma or serum was tested instead. Some small floating aggregates gradually developed over 4 days, and at day 7, some dendritic cells were evident. Although these cells exhibited potent T cell stimulatory capacity in the allo MLR, [Table 1, fourth row], the yield of dendritic cells was low, only  $0.5 \times 10^6$  cells per 40 ml of blood. As with FCS, the cells exhibited variable CD14 and CD32 expression and failed to express CD83 and p55. Thus human plasma was even less efficient than FCS in priming for mature dendritic cell development.

#### Requirement for conditioned medium

Two approaches were tested to increase dendritic cell yield and maturity in FCS and human plasma. A 1:10,000 dilution of Pansorbin was added to the cultures after 7 days of growth in GM-CSF and IL-4, and the cells were recultured for 3-4 days. Pansorbin increased the number

-34 -

of free floating dendritic cells as well as T cell stimulatory capacity in the allo MLR [Table 1, second and fifth rows]. However, there was a dramatic stimulation of syngeneic T cells possibly because of superantigens in the pansorbin.

In the second approach, we used conditioned medium [CM] derived from supernatants of T-depleted mononuclear cells that had been adhered to gamma globulin-coated plates or stimulated with pansorbin [see Methods]. On day 7 of culture, one half of the medium was replaced with this CM. At day 11, 4 days after the addition of CM, both the yield and enrichment of dendritic cells increased. [Table 1, rows 3 and 6]. There was also significant enhancement of allo stimulatory capacity, but without the increase in the syngeneic MLR seen with pansorbin. Thus we chose to use CM and 1% human plasma for all subsequent experiments. Plasma is preferable to use over serum since it is readily obtained during the Ficoll separation of blood.

In > 15 experiments, the overall effect of adding CM to plasma containing medium was to increase dendritic cell yields 3-10 fold [range of  $0.3 - 3 \times 10^6$  per 40 ml of blood] and enhance enrichment from 2-25% to 30-90%. The contaminating cells are primarily lymphocytes, B cells and residual T cells. CM generated cells were phenotypically and functionally stable in that they retained their dendritic cell features when cultured in the absence of cytokines [see below]. No substantial difference was noted whether 33% vs. 50% CM was used or whether the CM was replenished every 2 days.

Throughout the above cultures, we noted the presence of residual adherent cells, most likely macrophages, in both components of the culture, i.e., during priming for 6-7d with GM-CSF and IL-4, and during maturation for 3-4d with CM. We therefore modified the protocol further. At

-35 -

day 7 of cytokine treatment, we directly transferred the nonadherent cells to fresh plates. We then cultured the cells for 4 additional days (day 11) in the absence or presence of CM. As expected, GM-CSF/IL-4 primed cells that were cultured without CM were suboptimal stimulators in the MLR [Figure 1, closed circles]. In contrast, there was a substantial increase in stimulatory activity when cells received CM, [open squares]. ER- cells that were cultured in Teflon beakers without exogenous cytokines throughout the 11 days were poor APCs [open circles].

Phenotypic analysis of dendritic cells grown from normal peripheral blood with GM-CSF, IL-4 and Conditioned Medium

The dendritic cells that were induced by this new culture procedure displayed a typical dendritic cell phenotype [Figure 2] and morphology [Figure 3]. The cells expressed high levels of MHC class II and accessory molecules for T cell stimulation [CD54, CD58, CD40, CD80, CD86]. The more lineage restricted antigens such as CD3, CD19, CD16, CD14 were absent. CD1a, CD4, CD11c and CD45RO were also expressed. CD68 or macrosialin (Ramprasad et al. 1995; Rabinowitz and Gordon, 1991), was expressed in a perinuclear pattern. This pattern is distinct from that seen with macrophages, where staining is seen throughout the cytoplasm. Only rare cells stained with the anti-Lag mAb that identifies Birbeck granules in epidermal dendritic cells.

Two new markers, CD83 and p55, recently have been shown to be selectively expressed by the small subset of mature dendritic cells in cultured human blood (Zhou et al. 1992; Zhou and Tedder, 1995; Mosialos et al. 1995). Both markers were expressed only at low levels by cells that are primed with GM-CSF and IL-4, but were unregulated following culture in CM. CD83 is a member of the Ig superfamily, and p55 is an EBV-inducible actin bundling

35

-36 -

protein in B cells (Zhou et al. 1992; Zhou and Tedder, 1995; Mosialos et al. 1995).

Stability of dendritic cells generated in  
GM-CSF/IL-4 and CM

The stability of the dendritic cells generated at various points during the two step culture protocol described above was also ascertained. ER- cells were cultured with GM-CSF/IL-4 for 7 days and then analyzed for stimulatory activity in an allo MLR. Although these cells could stimulate allogeneic T cells [Table 2, first row], the phenotype was not typical of mature DCs [CD14+, moderate levels of HLA-DR, little to no CD83, and a diffuse staining for CD68; data not shown].

TABLE 2

<u>Treatment</u>				<u>Allogeneic MLR</u>		
Culture supplements on days:				<sup>3</sup> H-TdR Incorporation (cpm x 10 <sup>3</sup> ) at T:APC ratio of:		
<u>0-7</u>	<u>7-11</u>	<u>11-14</u>	<u>Day of MLR test</u>	<u>100:1</u>	<u>300:1</u>	<u>900:1</u>
GM+IL-4	-	-	7	161	145	81
"	no CM	-	11	203	146	25
"	+ CM	-	11	154	172	161
"	+ CM	no CM	14	144	170	189
"	+ CM	+ CM	14	178	207	165

Table 2: CM is required for optimal and stable  
immunostimulatory activity

ER- cells were cultured in GM/CSF and IL-4 for 7 days, washed and then analyzed immediately for stimulatory activity in an allo MLR [first row], or cultured for 4 more days in the absence or presence of CM [second and third rows]. At day 11, aliquots of dendritic cells were maintained in CM for three more days [row 4] or washed extensively and recultured in the absence of CM [row 5]. The same donor of allogeneic T cells was used throughout

-37 -

these experiments. Results are shown as  $^3\text{H}$ -TdR incorporation [cpm  $\times 10^3$ ] and are averages of triplicates.

Cells washed out of cytokines at day 7 and recultured for 4 more days in the absence of CM had a similar phenotype and a noticeable decline in T cell stimulatory function when compared with dendritic cells grown in GM-CSF/IL-4 and CM [compare rows 2 and 3, Table 2]. Cells were next washed out of CM at day 11 and cultured for three more days [day 14] alongside cells that were left in CM. The dendritic cells were identical with respect to their phenotype and stimulatory capacity in the allogeneic MLR [rows 4 and 5, Table 2]. Thus, the exposure to CM renders an apparent irreversible change that leads to functionally mature dendritic cells.

#### Attempts to identify the active cytokines in CM

The CM from blood cells that are cultured in human gamma globulin likely contain many monocyte-derived cytokines such as IL-1, IL-6 and  $\text{TNF}\alpha$ . However, none of these cytokines could replace the CM, and in fact each cytokine had no effect on increasing dendritic cell yield or function [Figure 4 and Table 3]. The cytokines IL-12 and IL-15 were also ineffective [Table 3].

-38 -

TABLE 3

5	Treatment		Allogeneic MLR		
	Culture supplements on day 7		<sup>3</sup> H-TdR Incorporation (cpm x 10 <sup>3</sup> ) at T:APC ratio of:		
	CM	Other	100:1	300:1	900:1
	+	goat IgG [4μg/ml]	147	117	46
	+	goat IgG [17μg/ml]	141	94	31
	+	α TNF-α [1μ/ml]	144	120	61
	+	α IL-6 [4μg/ml]	161	120	31
10	+	α IL-12	180	104	43
	+	α IL-1-α	152	90	33
	+	α IL-1-β	163	112	37
	+	cocktail [17μg/ml]	116	59	19
	-	TNF-α [10ng/ml]	79	38	11
15	-	IL-1-β	72	47	4
	-	IL-12 [500pg/ml]	55	21	4
	-	IL-15	46	12	4

**Table 3 : Attempts to identify the factors in CM that promote dendritic cell differentiation**

20 ER- cells were cultured in GM/CSF and IL-4 for 7 days, and then supplemented with CM for 4 days in the presence of neutralizing antibodies to a panel of cytokines. Goat IgG was the control. Cocktail refers to the combination of the various antibodies. Some cultures were supplemented with recombinant cytokines alone in the absence of CM. At day 11 of culture the cells were  
 25 evaluated for their ability to stimulate allogeneic T cells. Results are shown as <sup>3</sup>H-TdR incorporation [cpm x 10<sup>3</sup>] and are averages of triplicates.

We also added polyclonal neutralizing antibodies to cytokines to the CM. However, antibodies to TNFα, IL-1, and IL-6 did not block the capacity of CM to increase  
 30 dendritic cell yields and function [Figure 4 and Table 3].

#### Source of DC progenitors in PBMC

The progenitor cells that give rise to dendritic cells produced according to an embodiment of the invention  
 35



-39 -

[GM-CSF + IL-4 followed by CM] where characterized. To assess the adherence of progenitors to plastic, ER- cells were adhered for 1 hour in 6 well plastic plates [ $2 \times 10^6$  per well] after which nonadherent cells were removed and replated in new dishes. Both populations were cultured in GM-CSF and IL-4 for 7 days followed by CM for 4 days. Adherent ER- cells yielded the highest number and purity of dendritic cells [Table 4]

TABLE 4

DC progenitors are present in both adherent and non-adherent ER - fractions.

Cell population	DC Yield/40ml blood	% Enrichment	Allogeneic MLR		
			<sup>3</sup> H-TdR incorporation (cpm x 10 <sup>3</sup> ) at T:APC ratio of:		
			100:1	300:1	900:1
ER -	0.8 - 1.5 x 10 <sup>6</sup>	79 - 90	265	243	169
ER -, adherent	1.3 - 2.8 x 10 <sup>6</sup>	94 - 96	226	222	147
ER -, nonadherent	0.8 - 1.7 x 10 <sup>5</sup>	27 - 36	219	192	73

Table 4: Dendritic cell progenitors are present in both adherent and nonadherent ER- fractions

ER- cells were adhered to plastic for 1 hr. Both adherent and nonadherent cells were collected and cultured with GM-CSF and IL-4 for 7 days, followed by CM. At day 11, the yield and % enrichment of dendritic cells was determined. The T cell stimulatory function of the ER- populations analyzed is also shown. Results are representative of 2 experiments and were similar to cells derived from bulk ER- cells in terms of immunostimulatory capacity. ER- nonadherent cells also generated some potent APCs, but the yield and purity of dendritic cells was substantially less, primarily because of a large number of contaminating lymphocytes. Thus progenitors appear to reside in both adherent and nonadherent populations of blood mononuclear cells.

-40 -

ER- cells were stained with PE-conjugated anti-CD14 and the cells were sorted in a FACStar into CD14 high and CD14 low fractions. Negative small cells were excluded by gating. These were then cultured for 7 days with GM-CSF and IL-4 followed by 4 days in CM. CD14 high cells were also evaluated with or without prior treatment with ionizing irradiation [Gy 15 or 30]. The greatest yield, percent enrichment and T cell stimulatory activity were found in the unirradiated CD14 high population [Table 5 and Figure 5].

TABLE 5

DC progenitors are enriched in the CD14 hi population

Cell population	Yield/40ml blood	% Enrichment
CD14 hi	1.2 - 2 x 10 <sup>6</sup>	66 - 97
CD14 lo	0.7 - 0.8 x 10 <sup>6</sup>	10 - 39
CDF14 hi, XRT	1 - 1.6 x 10 <sup>6</sup>	72 - 77
CD14 hi + lo	1 - 1.3 x 10 <sup>6</sup>	57 - 73

[n = 2]

Table 5: Dendritic cell progenitors are enriched in the CD14 hi population

ER- cells were stained with CD14-FITC and sorted into CD14 hi and dim populations. Separated fractions, irradiated CD14 hi cells [Gy 15 and 30] and stained unsorted or combined cells [CD14 high + low] were cultured with GM-CSF/IL-4 followed by CM. The yields and % enrichment of dendritic cells is shown.

Interestingly, irradiated CD14+ high cells generated effective dendritic cells but the yield was 50% lower, and the percent enrichment was slightly lower, 77% vs. 97% for nonirradiated CD14+ cells. The CD14 low fraction also produced some typical dendritic cells, but the yield and

-41 -

enrichment were significantly lower than the CD14 high population, probably because there is a significant number of lymphocytes contaminating this fraction.

Collectively, our data suggest that dendritic cell progenitors are found in PBMC populations that are both plastic adherent and nonadherent, as well as CD14 dim and CD14 high. Furthermore, some of these progenitors are radioresistant.

10 Dendritic cells derived from non-proliferating progenitors induce strong anti-viral cytolytic T cell responses

Potent human CD8+ cytolytic T cell [CTL] responses to live replicating influenza A virus are generated when dendritic cells are the APCs (Bhardwaj et al. 1994). When pulsed with poorly replicating, heat-inactivated, influenza virus, dendritic cells induce equally strong CTL responses (Bender et al. 1995). Other APCs, e.g., macrophages lack this capacity. Since the heat inactivated virus is incapable of substantial new protein synthesis, only small amounts of viral protein are required to charge class I molecules on dendritic cells. Dendritic cells prepared as described above from non-proliferating progenitors in blood were tested for their capacity to elicit anti-viral CTL responses to live and heat-inactivated influenza virus [Table 6].

25

30

35

-42 -

TABLE 6

		Percent specific lysis of targets	
5	Infection of dendritic cells	Mo [-]	Mo [Flu]
	[-]	2.5	1.9
	Live Flu	1.3	60.4
10	Infection of dendritic cells	T2 [-]	T2 [Matrix peptide]
	[-]	0	10
	Heated Flu	30	77
15			

**Table 6:** Dendritic cells derived from non-proliferating progenitors induce strong anti-viral cytolytic T cell responses

20 Dendritic cells were uninfected or infected with live replicating or heat inactivated influenza virus [see methods] and cocultured with autologous HLA A2.1+ T cells [T:APC ratio = 30:1].

25 After 7 days, T cells were harvested and tested for cytolytic activity on chromium labeled syngeneic macrophages [mos] which were infected or uninfected with influenza virus [upper panel]. Lytic activity of T cells responding to dendritic cells infected with heat inactivated influenza virus was tested on chromium labeled T2 target cells [lower panel] that had been pulsed with 10 nM of HLA A2.1 restricted matrix peptide. E/T ratio in both experiments was 30:1.

30 Both forms of virus could be presented to autologous T cells to induce CTL responses. CD8+ T cells are the likely mediators, since they efficiently lysed TAP deficient T2 cells pulsed with the HLA A2.1 restricted influenza matrix peptide.

35

-43 -

Conditions for generating mature dendritic cells from progenitors in human plasma

The method of preparing mature dendritic cells according to this invention preferably uses two steps. Two steps are preferred because dendritic cells generated in either FCS or human plasma containing medium failed to develop and maintain a stable phenotype and function when removed from GM-CSF and IL-4. For example, these "putative" dendritic cells continued to express CD14 and CD32, which are typically absent from mature dendritic cells, expressed little or no CD83 and p55 and lost stimulatory function upon removal of GM-CSF/IL-4. This phenomenon has not been previously appreciated, since the cells are typically used without reculturing them in the absence of cytokines (Sallusto and Lanzavecchia, 1995; Sallusto and Lanzavecchia, 1994). Second, exposure to a dendritic cell maturation factor such as CM was an absolute requisite to induce the formation of large numbers of fully differentiated dendritic cells from the GM-CSF and IL-4 primed cultures. The CM-induced cells expressed a typical phenotype, with strong expression of a) antigen presenting MHC class I and II products and CD1a, b) several adhesions and co-stimulator molecules including CD40, CD54, CD58, CD80 and CD86, and c) two dendritic cell restricted molecules, CD83 and p55. This phenotype remained stable following removal of all growth factors. Moreover, the yield and T cell stimulating potency of the CM-treated cells were substantial. Up to  $3 \times 10^6$  dendritic cells were obtained from 40 ml of blood, which is 3 times more than the number of mature dendritic cells that are present in one unit (500 ml) of non treated blood cells. The cells induced strong allo MLRs even at DC:T ratios of 1:900. It is preferred to first prime the blood progenitors with GM-CSF and IL-4 for 6-7 days before

-44 -

adding the CM, since the latter seems to be less effective if added at day 0 [data not shown].

Dendritic cell derived from blood progenitors express a unique pattern of antigens

5 We find that CD83 and p55 are useful markers for the development of mature dendritic cells from blood progenitors. Both antigens are uniformly expressed at high levels in mature cells but are lacking in fresh blood monocytes and in progenitors that are primed in GM-CSF/IL-10 4 for 7 days but are not matured in CM. Another useful marker to distinguish dendritic cells from monocytes is CD68 [termed macrosialin in the mouse], a member of the lamp-1 family. This antigen is detected in a perinuclear area in dendritic cells, whereas much of the cytoplasm 15 stains for CD68 in monocytes, as has been emphasized by Gordon and others (Rabinowitz and Gordon, 1991; Ramprasad et al. 1995).

EXAMPLE 2

20 Materials And Methods  
Culture medium

The following culture media were used: RPMI 1640 [Biological Industries, Kibbutz Beit Haemek Israel]; X-VIVO10, 15, and 20™ [Bio-Whittaker, Walkersville, MD]; 25 HybriCare and AIM-V™ [GIBCO-BRL, Gaithersburg, MD]. They were supplemented with 50µg/ml of gentamicin. Sources of serum were FCS [10%, Biological Industries or Seromed-Biochrom, Berlin, Germany], and human plasma and serum [0.5, 1, 5, and 10%] from either adults [PAA Laboratories 30 GmbH, Lin, Austria, blood bank or autologous] or from umbilical cord blood [kindly provided by the Department of Gynecology and Obstetrics]. FSC and, in some experiments only, human sera/plasma was heat-inactivated at 56°C for 30 minutes.

35

-45 -

Cytokines and anti-cytokines

Recombinant human GM-CSF was generously provided by Dr. E. Liehi, Sandoz Ges.m.b.H., Vienna, Austria [specific activity  $5.9 \times 10^6$  U/mg]. Alternatively, human GM-CSF prepared for therapeutical purposes was used [Leukomax™, Sandoz, Basel, Switzerland, specific activity  $1.1 \times 10^7$  U/mg]. Recombinant human IL-4 was a gift of Dr. M.B. Widmer, Immunex Corp., Seattle, WA [specific activity  $5 \times 10^7$  U/mg] and Dr. E. Liehl, Sandoz, TNF- $\alpha$  was supplied by Dr. G.R. Adolf [Bender, Vienna, Austria; specific activity  $6 \times 10^7$  U/mg]. IL-12 was provided by the Genetics Institute, Boston, MA [specific activity  $3.6 \times 10^8$  U/mg]. Human IL-1 $\beta$  [specific activity  $5 \times 10^8$  U/mg] and simian IL-15 [specific activity  $1 \times 10^7$  U/mg] were purchased from Genzyme Corp., Cambridge, MA human stem cell factor [specific activity  $0.5-1.0 \times 10^6$  U/mg] from R&D Systems [MO]. Human IL-15 and a neutralizing rabbit anti-human IL-15 antibody were from PeproTech, London, UK. Human IL-13 was a gift of Dr. A. Minty, Sanofi, Labège, France and was also obtained from Pharmingen, San Diego, CA [specific activity  $1 \times 10^6$  U/mg].

Initial processing of human blood

Peripheral blood was obtained from the local blood bank as standard buffy coat preparations and from normal donors. Preservative-free heparin [200 I.U./ml blood; Novo Nordisk A/S, Bagsvaerd, Denmark] was used to prevent clotting. PBMC were isolated by centrifugation in Lymphoprep [1.077 g/ml; Nycomed Pharma AS, Oslo, Norway]. In order to minimize contamination of PBMC with platelets the Lymphoprep centrifugation [200 x g/room temperature] was interrupted after 20 minutes. The top 20-25 ml containing most of the platelets were carefully removed and centrifugation was resumed [20 minutes/460 x g/room temperature]. PBMC were then depleted of T and B cells by

-46 -

means of an immunomagnetic technique. Dynabeads [Dyna-  
Oslo, Norway] M-450 Pan-B/CD 19 and M-450 Pan-T/CD2 were  
washed four times with PBS containing 1% bovine serum  
albumin or human plasma. PBMC [ $30-50 \times 10^6/\text{ml}$ ] were  
5 incubated with the magnetic beads at a ratio of 1:1:1 at  
4°C for 20 min. using Dynal mixer. Lymphocytes were then  
depleted by means of a Dynal magnet [1-2 minutes].  
Magnet-non-adherent cells were harvested and washed. This  
depletion step was repeated once. In some experiments an  
10 indirect immunomagnetic approach was used. PBMC were  
labeled with anti-T and B cell mAb's [hybridoma  
supernatant of mAb OKT-11/CD2 and ascitic fluid of mAb  
MEM-97/CD20; gift of Dr. v. Horejsi, Praha, Czech  
Republic], washed three times and incubated with sheep  
15 anti-mouse Ig-coupled magnetic beads [SAM-M450; Dynal] at  
a ratio of 3 beads to one cell. Magnetic separation was  
achieved as described above. In addition, lymphocyte-  
depleted PBMC were obtained from cancer patients [after  
informed consent] in complete remission during  
20 hematopoietic recovery after high-dose consolidation  
chemotherapy and subcutaneous daily administration of  
300 µg G-CSF [Neupogen™, Hoffmann-La-Roche, Basel,  
Switzerland]. Depletion of PBMC from CD34+ cells was  
achieved either by means of M-450/CD34 magnetic beads  
[Dynal] or by passing cells through an immunoaffinity  
25 anti-CD34 column [CellPro Inc., Bothell, WA] that is  
clinically used for transplantation purposes. PBMC from  
normal donors treated with G-CSF in the same manner were  
also used. This study was approved by the Ethical  
Committee of the Medical Faculty, University of Innsbruck.

30

#### Culture technique

Lymphocyte-depleted PBMC were planted in 6-well  
tissue culture plates at a density of  $2 \times 10^6$  cells/well  
in 3 ml of complete culture medium. GM-CSF and IL-4 were

35



-47 -

added at final concentrations of 800 and 1000 U/ml, respectively. Cultures were fed every other day [days 2, 4 and 6] by removing 1 ml of the medium and adding back 1.5 ml fresh medium with cytokines [1600 U/ml GM-CSF and 1000 U/ml IL-4, resulting in final concentrations of 800 and 500 U/ml, respectively]. On day 7, non-adherent cells were either harvested and analyzed or transferred to new 6 well plates and cultured further in the presence or absence of maturation stimuli [see below]. GM-CSF and IL-4 were present during the culture period from day 7 to day 10-11.

#### Stimuli for dendritic cell maturation

SACS [fixed *Staphylococcus aureus* Cowan 1 strain, 2.01 mg/ml Ig-binding capacity, Pansorbin cells, Cat No. 507861] was purchased from CalBiochem Co., La Jolla, CA. It was added to dendritic cell cultures at a final dilution of 1:10000. Conditioned medium was produced by Ig-adherent PBMC. To this end, bacteriologic plates [100 mm. Cat. No. 1029, Falcon, Oxnard, CA] were incubated with a solution of human Ig [10 mg/ml, therapeutically used "Immunoglobulin i.v. Biochemie", Biochemie Ges.m.b.H., Vienna, Austria, subsidiary of Sandoz] in PBS for 1 minute at room temperature. After three rinses with PBS  $5 \times 10^7$  PBMC were put into one Ig-coated bacteriologic plate for 1 hour at 37°C in 8 ml of complete culture medium. Non-adherent cells were rinsed away and adherent cells were cultured in fresh complete medium at 37°C for 24 h.

Complete media consisted of the different media and serum supplements mentioned above. Conditioned media were sterile filtered and stored at +4°C for one week at the longest or frozen at -20°C. In some cases, conditioned medium was prepared from cells stimulated with SCS instead of human Ig. Conditioned media were routinely used at 25%

35

-48 -

vol/vol. Increasing this concentration up to 50% was not appreciably better.

### Flow cytometry and immunohistochemistry

Primary antibodies used for flow cytometric and immunohistochemical analyses are listed in Table 7.

TABLE 7

Antibodies used for immunohistochemistry					
	Specificity	Clone/Name	Ig class	Source	Reference
10	HLA-DR/DQ	9.3F10/HB 180	mouse IgG2a	ATCC <sup>a</sup>	
	HLA-DR	L243/HB55	mouse IgG2a	ATCC	
	HLA-DR-FITC	L243	mouse IgG2a	BDIS <sup>b</sup>	
	CD1a	OKT-6/CLR8020	mouse IgG1	ATCC	
	CD1a-FITC	OKT-6	mouse IgG1	Ortho <sup>c</sup>	
15	CD15s	CSLEX1	mouse IgM	BDIS	
	CD40	G-28.5	mouse IgG1	E.A. Clark <sup>d</sup>	
	CD44v9	VFF16	mouse IgG	P.Herrlich <sup>e</sup>	
	CD45RA	4G10	mouse IgG2a	R.M. Steinman <sup>f</sup>	
	CD45RO	UCHL10	mouse IgG2a	DAKO <sup>g</sup>	
20	CD68	EBM11	mouse IgG1	DAKO	
	CD71	RPN.511	mouse IgG1	Amersham <sup>h</sup>	
	CD80	L307.4	mouse IgG1	BDIS	
	CD83	HB-15a	mouse IgG2b	T.F. Tedder <sup>i</sup>	(Zhou et al. 1992)
	CD86	IT2.2	mouse IgG2b	Pharmingen <sup>j</sup>	
25	CD115/c-fms	Ab-1	rat IgG1	Oncogene	Ashmun et al., 1989
	CSF-1-receptor	(clone 2-4A5-4)		Science <sup>k</sup>	
		Ab-2	rat IgG2b		
		(clone 3-4A4-E4)			
	Birbeck granules	Lag	mouse IgG1	S. Imamura <sup>l</sup>	(Kashihara et al. 1986)
30	Proliferation-associated	Ki-67	mouse IgG1	DAKO	

35

TABLE 7

## Antibodies used for immunohistochemistry

	• American Type Culture Collection, Rockville, MD;
	• Becton-Dickinson Immunocytometry Systems, Mountain View, CA;
5	• Ortho, Raritan, NJ;
	• Seattle, WA;
	• Kernforschungszentrum, Karlsruhe, Germany;
	• The Rockefeller University, New York, NY;
	• Roskilde, Denmark;
	• Amersham, UK;
	• Duke University, Durham, NC;
	• San Diego, CA;
10	• Cambridge, MA;
	• Kyoto University, Kyoto, Japan

Secondary antibody was biotinylated anti-mouse or anti-rat Ig's followed by FITC-or PE-conjugated streptavidin [Amersham International, Amersham, UK]. Dead cells were gated out on the basis of their fluorescence with propidium iodide or by their light scatter properties. Analyses were done on a FACScan instrument [Becton-Dickinson, Mountain View, CA]. Cytospins were prepared with Cytospin 2 centrifuge [Shandon, Inc., Pittsburgh, PA]. Slides were acetone-fixed for 5 min. at room temperature and then incubated with primary mAbs for 15 minutes followed by biotinylated sheep anti-mouse Ig [1:200; Amersham] and Texas Red-conjugated streptavidin [1:50, Amersham]. For double labeling purposes this staining sequence was extended with a blocking step [100 µg/ml mouse gamma globulin] and FITC-conjugated anti-HLA-DR [Becton-Dickinson]. Slides were mounted in Vectacshield™ [Vector Laboratories, Burlingame, CA]

### 30 Allogenic MLR and processing/presentation of soluble antigen

To test for T cell stimulatory function, the APCs were irradiated with 30 Gy from a Cs source and added in graded doses as simulators for  $2 \times 10^5$  purified [nylon

wool-non adherent; anti MHC class II-panned], allogeneic T cells in 96 well flat bottomed plates [Falcon]. For some experiments T cells were purified from umbilical cord blood in order to obtain naive T cells. Proliferation was determined on days 4-6 with the addition of 4  $\mu$ Ci-148 KBq/ml of [ $^3$ H]TdR [specific activity 247.9 GBq/mmol=6.7 Ci/mmol; New England Nuclear, Boston, MA] for 12-16h to triplicate wells [mean cpm]. Processing and presentation of tetanus toxoid protein [Connaught, Willowdale, Ontario, Canada] was measured with a HLA-DP4- restricted tetanus-peptide-specific T cell clone [AS11.15] that was a gift of Dr. A. Lanzavecchia, Basel Switzerland (Lanzavecchia, 1985). Graded doses of irradiated dendritic cells were co-cultured with  $1.5 \times 10^4$  clone cells in the presence or absence of tetanus toxoid [5 and 1  $\mu$ g/ml]. [ $^3$ H] TdR was added from d2 to d3.

To generate cytotoxic T cells, the approach described by (De Bruijn et al. 1992) was used with modifications. Irradiated [12.5 Gy] dendritic cells as simulators were co-cultured with autologous PBMC or with CD8+ T lymphocytes [1:25] as responders in the presence of 100-200  $\mu$ M peptide MP 58-66 of influenza matrix protein [MedProbe, Oslo, Norway] in 2ml RPMI/10% human serum in wells of a 24-cell tissue culture plate. After one week the cultures were fed with 40% Lymphocult [Biotest, Dreieich, Germany]. The lytolytic potential of outgrowing T cells was assessed two to four weeks after the start of the cultures in a standard Chromium-release cytotoxicity assay using as target PHA-stimulated autologous T cell blasts, that had been incubated with different concentrations of the peptide. Results are expressed as % specific lysis. CD8+ T lymphocytes were prepared by indirect immunomagnetic depletion of CD4+ T cells using mAb VIT-4/anti-CD4 [gift of Dr. W. Knapp, Vienna, Austria] followed by sheep anti-mouse IgG magnetic beads [Dynal].

o Additional reagents

Latex beads [0.5% vol/vol; 2  $\mu$ m diameter] for phagocytosis experiments were purchased from Seradyn, Indianapolis, IN. They were added to the cell cultures at a final dilution of 1:20. Staphylococcal enterotoxin A [superantigen SEA] was from Sigma Chemical Corp. St. Louis, MO. It was used at a final concentration of 25  $\mu$ g/ml. LPS [Sigma] was added to some experiments at a final concentration of 100 ng/ml.

10 RESULTS

Identification of culture conditions allowing full maturation of dendritic cells

Dendritic cells grown in GM-CSF and IL-4 are not fully mature. Lymphocytes were depleted from PBMC by means of immunomagnetic beads. Use of the immunomagnetic beads allowed for the omission of a sensing step two hours after plating. Dendritic cells developed from these lymphocyte-depleted PBMC in the same way as from whole PBMC populations.

Dendritic cells, that had been washed out of GM-CSF and IL-4 and were replated in culture medium without cytokines for an additional three days [days 7-10], lost their characteristic morphology and re-adhered. They assumed the typical shapes of macrophages [Figure 6A]. In addition, dendritic cells on d7 of culture expressed the macrophage marker CD 115 [c-fms/CSF-1 receptor] although their CD14 expression was very low or absent. In contrast, they did not express CD83, a marker for mature dendritic cells (Zhou and Tedder, 1995) [Figure 7]. The staining pattern of a well defined population of mature dendritic cells, namely cutaneous dendritic cells obtained by emigration from skin organ cultures (Pope et al. 1995) was inverse: CD115/CD83+ [not shown]. From this we concluded that dendritic cells grown in GM-CSF and IL-4

-52 -

- o were not fully and irreversibly mature although they did express some maturation markers such as high levels of MHC class II, furthermore CD40, 54, 58, 80 and a strong T cell stimulatory capacity (Romani et al. 1994; Sallusto and Lanzavecchia, 1994).

5 Determination of stimuli for full maturation of dendritic cells

- We tested whether full maturation could be brought about by adding cytokines to the cultures. Supplementation of GM-CSF and IL-4 containing culture medium with IL-18 [50U/ml] and/or TNF- $\alpha$  [50U/ml] either from day 0 through to d7 or as a 24-hour pulse from day 0 to day 1 or day 6 to day 7 increased immunostimulatory capacity as expected but did not lead to a stable phenotype of mature dendritic cells as defined above. Addition of these cytokines after the initial 7-day culture and transfer of the cells to fresh culture wells [i.e., from day 7-10] gave the same result [Figure 8]. Similar findings were made for the cytokines IL-12 [1 ng/ml], stem cell factor/c-kit ligand [yyU/ml] and IL-15 [20ng/ml].
- SACS [fixed *Staphylococcus aureus*], a potent natural stimulator of cytokine secretion was also tested. When a 1:10,000 dilution of SACS was added to the cultures from day 7 to day 10 or 11 of culture, the numbers of free floating dendritic cells increased markedly. Moreover, these cells displayed a more pronounced morphology with highly motile cytoplasmic processes ["veils"]. The cells also did not re-adhere when replated without cytokines. T cell stimulatory capacity in the allogeneic MLR was superior to dendritic cells that had been cultured without SACS [Figure 8]. However, there was a dramatic stimulation of syngeneic T cells possibly because of superantigens in the SACS preparation. Addition of superantigen to the cultures between days 7 to 10 did not

35

- o bring about maturation of dendritic cells. However, supernatants of SACS-stimulated adherent PBMC [i.e., conditioned media] had the same effect as SACS itself.

Conditioned medium [CM] was produced in a way that could potentially be of clinical use and avoid the involvement of bacterial products such as SACS.

5 Therefore, we derived supernatants from whole or T-depleted mononuclear cells that had been stimulated by adherence to Ig-coated plates. On day 6 to 8 [mostly 7] of culture dendritic cells were transferred to fresh wells and 25% vol/vol of CM was added. Three to four days later

10 the cells were analyzed. There were almost no adherent cells left on the bottom of the wells. By phase contrast microscopy revealed that the floating cells had many motile veils, much like SACS-treated cells [Figure 9]. In the hemocytometer the cells appeared "hairy" like cultured

15 epidermal Langerhans cells or cutaneous emigrant dendritic cells. This is in contrast to either 7 day or 10 day cultures without CM, where the cells are large with irregular outlines, but the number of long processes and veils is low. Cultured without cytokines the CM-exposed

20 cells remained stably non-adherent over an observation period of 3 days [Figure 6B-D]. They now expressed CD83 and were highly stimulatory for resting T cells [see below]. Therefore, dendritic cells generated in the presence of monocyte CM qualified as fully mature

25 dendritic cells. In >15 experiments the yields of mature, CD83-expressing dendritic cells per 40 ml of blood ranged between 1 and  $4 \times 10^6$  in FCS-containing media. Maturation occurred likewise when IL-4 was replaced with IL-13 [20 ng/ml].

30 To ascertain whether the activity of CM is due to IL-15, a neutralizing antibody was added to CM during the maturation phase of culture. Maturation was not prevented or impaired. We also measured whether CM contained

- bioactive IL-15. IL-2-responsive CTLL-2 cells also proliferated vigorously in response to both human and simian IL-15 in concentrations as low as 5 pg/ml. No response was seen when CM were tested in this assay [n=2] indicating that IL-15 is present only in minute quantities if at all [Table 8].

TABLE 8

Test for IL-15 in the monocyte conditioned medium

Additions to the CTLL bioassay for IL-15	no antibody	anti-IL-5 1:100	anti-IL-15 1:200	anti-IL-15 1:100 + excess IL-15
no cytokines	0.2			
IL-15 [500 pg/ml]	152.8	148.8	161.1	149.2
IL-15 [50 pg/ml]	151.1	145.3	147.0	144.2
IL-15 [5 pg/ml]	45.8	0.9	0.8	156.9
monocyte conditioned medium [25% vol/vol]	0.6	2.4	1.1	155.0

Table 8: IL-15-dependent proliferation of CTLL-2 cells was determined by measuring  $^3\text{H}$ -TdR incorporation during the final 6 hours of a 36 hour culture [ $6 \times 10^3$  CTLL cells per well]. Values are means of triplicate wells and are given in cpm  $\times 10^3$ . Excess IL-15 was added at a final concentration of 1 ng/ml.

#### Development of FCS-free media

Having established the conditions for the full maturation of dendritic cells it was desirable to develop a system without fetal calf serum. Simply replacing FCS with human pool sera resulted in little or no development of dendritic cells. We observed increased adherence of cells to the culture surfaces. Serum, plasma and serum from umbilical cord blood were tested over a concentration



0 range from 0.5 to 10%. We also compared autologous  
sera/plasma versus pooled sera/plasma from the blood bank  
or commercially available reagents. The effects of heat-  
inactivation were tested. These variables were combined  
with different culture media such as RPM1640, X-VIVO, AIM-  
5 V, Hybricare, and Iscove's. These reagents were used for  
the culture of both progenitor cells and CM-producing  
adherent PBMC. The primary read-out system was morphology  
under the inverted phase contrast microscope and the yield  
of mature, stable dendritic cells. Cultures in RPMI  
10 supplemented with 10% FCS were always run in parallel as a  
positive control. RPMI1640 containing 1% autologous, not  
heat-inactivated human plasma was observed to be optional.  
In 12 standardized experiments the yield ranged between  
0,8 to  $3,3 \times 10^6$  CD83+ dendritic cells from 40 ml of  
15 blood. Enrichment of CD83+ cells was 30 to 80%.

#### Properties of mature dendritic cells

Phenotype. FACS analyses were performed before and  
three days after exposure to CM. Alternatively, dendritic  
cells were cultured from day 7 to day 10 in the presence  
20 or absence of CM and then compared to each other. Figure  
7 shows expression of the key markers: CD83 is induced,  
CD86 is enhanced, and CD115 is lost. This pattern applies  
also to dendritic cells cultured in clinically approved  
media and derived from PBMC that have been depleted from  
25 lymphocytes by immunomagnetic beads or sheep erythrocyte  
resetting. CD14 is low or absent and CD1 a shows some  
reduction in expression levels upon maturation. Further  
CM-induced phenotypical changes include the down-  
regulation of CD32 and CD45RA and the upregulation of  
30 CD45RO. Substantial levels of CD4 are expressed on both  
immature and mature dendritic cells. Staining of  
cytocentrifuge smears [not shown] also reflected the  
maturation event that takes place under the influence of

CM. Dendritic cells on day 7 of culture are stained throughout the whole cytoplasm in a pattern typical for monocytes or macrophages. After three days of culture in the presence of CM the immunostained cellular structures become concentrated in a dull spot near the nucleus. This pattern has been described for mature dendritic cells. MAb's Lag and Ki-67, specific for Birbeck granules of epidermal dendritic cells [i.e., Langerhans cells] and a proliferation-associated antigen, respectively, stained only rare dendritic cells.

Function. One of the mechanisms by which dendritic cells take up antigens is phagocytosis. This function is down-regulated upon culture of dendritic cells (Reis e Sousa et al. 1993). We tested the phagocytic capacity of dendritic cells before and after exposure to CM. Immature dendritic cells readily took up 2 $\mu$ m latex beads; mature dendritic cells after three days of culture in CM phagocytosed only very few particles [Figure 10].

Maturation in conditioned medium enhanced the stimulatory capacity for resting T cells markedly (Figures 11A, 11B). Cord T cells, i.e., naive T cells were also efficiently stimulated by mature dendritic cells [Figure 11C]. When we tested for the capacity to process and present a soluble native protein antigen [tetanus toxoid] using a tetanus peptide-specific T cell clone a reciprocal pattern emerged. Dendritic cells that had been cultured in the presence of CM were less efficient than those cultured in the absence of CM [Figure 11D] indicating maturation (Sallusto and Lanzavecchia, 1994; Romani et al. 1989a). Mature dendritic cells, when pulsed with a dominant influence matrix peptide that is presented on HLA-A2.1 were able to elicit peptide-specific, MHC class I-restricted, lytic T cells lines from autologous PBMC or populations of CD+ T cells [Figure 11E, F].

° Modifications of the method with regard  
to clinical applicability

With regard to the clinical application of cultured dendritic cells we tested whether the method would work in the complete absence of xenoproteins and with culture media that are approved for clinical use. Comparative  
5 data showed that depletion of lymphocytes [T and B] with immunomagnetic beads was equivalent to the commonly used method of rosetting with neuraminidase-treated sheep erythrocytes discussed above. Similar numbers of stimulatory dendritic cells were obtained. Anti-B and T  
10 cell mouse mAb's and immunomagnetic sheep anti-mouse Ig beads, both approved for clinical use are available from Baxter Healthcare Corp. [Glendale, CA]. Culture media for clinical use are optimized for the growth of cell types other than dendritic cells [e.g. tumor-infiltrating  
15 lymphocytes]. Yet, in more than 15 experiments mature dendritic cells could consistently be procured. When X-VIVO 20 medium supplemented with 1% autologous human plasma was used for both culturing the cells and producing CM, substantial numbers of stimulatory dendritic cells  
20 could be obtained. Yields and percentages of enrichment were, however, only about half of what could be achieved with RPMI 1640 medium [n=5]. More adherent cells were observed in these cultures. Pilot experiments with AIM-V medium gave similar results. X-VIVO 10 and X-VIVO 15  
25 media yielded unsatisfactory results.

With regard to the possibility that only small volumes of blood may be available in certain clinical settings we determined that pretreatment of donors with G-CSF increased dendritic cell yields. Blood from G-CSF-  
30 treated cancer patients and normal individuals gave up to 6-fold yields of mature [i.e., CD83+] dendritic cells. This increase was not only due to an increased frequency of CD34+ precursor cells in the blood because similar

-58 -

numbers of immunostimulatory mature dendritic cells could be grown both from CD34-containing and CD34-depleted PBMC starting populations in response to GM-CSF, IL-4 and CM [Figure 12].

We also tested whether cryopreservation of dendritic cells precursors and their progeny is possible. Indeed, using standard freezing procedures dendritic cells could be generated from thawed populations of PBMC. Although not investigated systematically, it is preferred to freeze the starting population of PBMC or immature dendritic cells on day 7 of culture [i.e.], the day of transfer into CM-containing medium rather than mature cells on day 10/11 of culture.

The criteria used initially to define fully mature dendritic cells included: morphology, lack of adherence to plastic, and select surface markers. Dendritic cells grown for six days in GM-CSF and IL-4, i.e., immature dendritic cells did not express CD83 but did express CD115 [CSF-1 receptor/M-CSF receptor]. When they were replated without cytokines [and without CM] they adhered to the bottom of the culture wells. After a three day culture in the presence of CM they became CD83+/CD115- and did not become adherent upon withdrawal of cytokines. They also maintained their pronounced cytoplasmic, motile "veils". These results are consistent with the production of fully and stably mature dendritic cells. Only SACS, conditioned media from SACS-stimulated adherent PBMC, and conditioned media from monocytes stimulated by adherence on Ig were able to make dendritic cells reach full maturation. When we tried to replace CM with cytokines we observed no effects or some increase in T cell stimulatory capacity [with IL-1 $\beta$ , and/or TNF- $\alpha$  or IL-15]. Full maturation was not achieved, however. In as shown in Example 1, neutralizing antibodies to IL-1 $\alpha$  and  $\beta$ , IL-6, TNF- $\alpha$ , IL-12, alone or in combination, do not block the activity of

35

- o the monocyte CM. CM may contain either a unique cocktail of cytokines that is difficult or impossible to reproduce experimentally or new cytokines.

### EXAMPLE 3

#### 5 Clinical Use of Dendritic Cells Primed With Influenza or HIV Antigens

In this Example, dendritic cells prepared as described in Example 1 are primed with peptides that are presented on MHC class I molecules to prime CD8+ T cells. We have used the immunodominant influenza matrix peptide, 10 GILGFVFTL, that is recognized by CTLs from HLA-2.1 individuals, and we will in addition use HIV-1 peptides [pol ILKEPVHC-V; gag SLYNTVATL] that are recognized by CTLs from HIV-1 infected individuals. The efficacy of the peptide pulse can be monitored by applying an aliquot of 15 the DCs to autologous T cells in culture and measuring the killer cell response.

Most humans are primed to influenza antigens, so that the recall CTL response to this virus provides an optimal and internal control to determine if autologous 20 DCs, pulsed ex vivo with viral antigens, can boost the CTL response in vivo.

In this protocol, HIV-1 infected individuals are treated with CD4+ T cell counts of 300-400/u1. These individuals have low numbers of HIV-1 specific, killer 25 cell precursors. Asymptomatic, HIV-1 infected, adult volunteers of both sexes will be recruited. The donors must be HLA-A2.1 positive, as will be determined by initial typing with a monoclonal to A2.1. Donors who have relatively low and high recall responses to influenza will be included. The therapeutic goal is to increase the 30 numbers of these killers in order to reduce the cell-cell spread of HIV-1. The killer cell response to influenza is an internal control, since it should be intact in HIV-1 infected patients.

-60 -

- o Pairs of HIV-1 peptides are used so that the response to dendritic cells administered by two routes, intradermal [i.d.] and subcutaneous [s.c.] can be compared in the same patient. The influenza peptide will serve as the common denominator for each route. The peptides are made according to Good Laboratory Practices [GLP] at Sloan Kettering Institute, where peptides have been synthesized previously for administration in humans.

The Clinical Protocol is as follows:

- 10 a. Take a 50-60 ml blood sample in heparin by venipuncture.
- b. Isolate the mononuclear cells on Ficoll Hypaque, and save the autologous plasma.
- 15 c. Separate the mononuclear cells into T cell enriched and T cell depleted fractions by sheep erythrocyte rosetting, or by depleting lymphocytes with GMP-grade DYNAL beads coated with monoclonal antibodies to B and T cells. The T cells are frozen for tissue culture tests for antigen presentation to cytolytic T lymphocytes [CTLs, below].
- 20 d. Culture a fraction of the mononuclear cells on culture plates that have been coated with human Ig [Sandoz] in an RPMI-1640 culture medium supplemented with antibiotics [penicillin and streptomycin]. This stimulates the cells to form a conditioned medium that
- 25 fosters DC maturation.
- e. Culture the bulk of the T-depleted mononuclear cells for 6 days in RPMI-1640 supplemented with antibiotics, 1% autologous plasma, 1000 U/ml rHuGM-CSF, and 1000 U/ml rHuIL-4. GMP grade cytokines are provided by
- 30 Schering-Plough.
- f. After 6 days, transfer the cells to fresh medium supplemented with 30-50% v/v autologous monocyte conditioned medium plus 1 uM of influenza matrix peptide,

-61 -

o. GILGFVFTL. One aliquot of the DCs is pulsed with the HIV-1 gag peptide and the other with the polymerase peptide. At day 11, when the DCs have acquired their typical stellate shapes, the DCs are harvested, washed, counted, and readied for injection by s.c. and i.d. routes. Prior to the injection, the patients are typed to be sure that they have killer cell precursors specific for the above peptides.

g. The DCs are injected in the forearm. On one side the cells are given in a total of 0.5ml in 3 intradermal sites, while on the other side, the cells are given in 0.5 ml in 3 subcutaneous sites. The purpose of the two routes is to compare the efficacy of boosting to the influenza matrix peptide.

h. 1 week and 1 month after injection, 25 ml of blood is taken to evaluate the induction of immune responses. T cell responsiveness to the peptide is measured by antigen-induced proliferation and release of IFN-gamma in culture. The results are compared to preinjection values which are obtained 0 and 1 month prior to injection. The first injection utilizes DCs that are not pulsed with peptide. T cell responsiveness to peptide will be measured by enumerating CTL precursor frequencies [CTLp] for the peptide-coated, HLA-A2.1 "T2" cell line.

i. If priming is not detected at 1 month, a second dose of antigen pulsed DCs is be administered. If priming is detected at 1 month, the immune status is reexamined at 3 months and 6 months to determine the longevity of immune memory.

While we have hereinbefore described a number of embodiments of this invention, it is apparent that the basic constructions can be altered to provide other embodiments which utilize the methods and compositions of the invention. Therefore, it will be appreciated that the scope of this invention is defined by the claims appended

35

-62 -

- hereto rather than by the specific embodiments which have presented hereinbefore by way of example.

5

10

15

20

25

30

35



o Bibliography

Alijagic, S., Moller, P., Artuc, M., Jorgovsky, K., Czarnetzki, B.M. and Schadendorf, D. (1995). Dendritic cells generated from peripheral blood transfected with human tyrosinase induce specific T cell activation. Eur. J. Immunol. 25, 3100.

5 Ashuman, R.A., Look, A.T., Roberts, W.M., roussel, M.F., Seremetis, S., Ohtsuka, M., and Sherr, C.J. (1989). Monoclonal antibodies to the human CSF-1 receptor (c-fms protooncogene product) detect epitopes on normal mononuclear phagocytes and on human myeloid leukemic  
10 blasts. Blood 73, 827-837.

Austyn, J.M. (1987) Lymphoid dendritic cells. Immunol. 62, 161.

Bakker, A.B.H., Marland, G., De Boer, A.J., Huijbens, R.J.F., Danen, E.H.J., Adema, G.J. and Figdor, C.G. (1995) Generation of antimelanoma cytotoxic T  
15 lymphocytes from healthy donors after presentation of melanoma-associated antigen-derived eptiopes by dendritic cells in vitro. Cancer Res. 55, 5330.

Bender, A., Bui, L.K., Feldman, M.A.V., Larsson, M. and Bhardwaj, N. (1995) Inactivated influenza virus, when presented on dendritic cells, elicits human CD8<sup>+</sup> cytolytic T cell responses. J. Exp. Med. 182, 1663.

Bhardwaj, N., Lau, L., Rivellis, M. and Steinman, R.M. (1988) Interleukin-1 production by mononuclear cells  
25 from rheumatoid synovial effusions. Cell. Immunol. 114, 405.

Bhardwaj, N., Bender, A., Gonzalez, N., Bui, L.K., Garrett, M.C. and Steinman, R.M. (1994) Influenza virus-infected dendritic cells stimulate strong  
30 proliferative and cytolytic responses from human CD8<sup>+</sup> T cells. J. Clin. Invest. 94, 797.

Boon, T., Cerottini, J.-C., Van den Eynde, B., Van der Bruggen, P. and Van Pel, A. (1994). tumor antigens recognized by T lymphocytes. *Annu. Rev. Immunol.* 12, 337.

Carr, C.M. and Kim, P.S. (1993) A spring-loaded mechanism for the conformational change of influenza hemagglutinin. *Cell* 73, 823.

Caux, C., Dezutter-Dambuyant, C., Schmitt, D. and Banchereau, J. (1992) GM-CSF and TNF-alpha cooperate in the generation of dendritic Langerhans cells. *Nature* 360, 258.

Caux, C., Massacrier, C., Vanbervliet, B., Dubois, B., Van Kooten, C., Durand, I. and Banchereau, J. (1994) Activation of human dendritic cells through CD40 cross-linking. *J. Exp. Med.* 180, 1263.

Caux, C., Liu, Y.-J. and Banchereau, J. (1995a) Recent advances in the study of dendritic cells and follicular dendritic cells. *Immunol. Today* 16, 2.

Caux, C., Massacrier, C., Dezutter-Dambuyant, C., Vanbervliet, B., Jacquet, C., Schmitt, D. and Banchereau, J. (1995b) Human dendritic Langerhans cells generated in vitro from CD34<sup>+</sup> progenitors can prime naive CD4<sup>+</sup> T cells and process soluble antigen. *J. Immunol.* 155, 5427.

Crowley, M., Inaba, K. and Steinman, R.M. (1990) Dendritic cells are the principal cells in mouse spleen bearing immunogenic fragments of foreign proteins. *J. Exp. Med.* 172, 383.

De Bruijin, M.L.H., Nieland, J.D., Schumacher, T.N.M., Ploegh, H.L., Kast, W.M. and Melief, C.J.M. (1992) Mechanisms of induction of primary virus-specific cytotoxic T lymphocyte responses. *Eur. J. Immunol.* 22, 3013.

Egner, W. and Hart, D.N.J. (1995). The phenotype of freshly isolated and cultured human bone marrow allositulatory cells: Possible heterogeneity in bone marrow dendritic cell populations. *Immunology* 85, 611.

0 Fossuam, S. (1989a) Lymph-borne dendritic leucocytes do not recirculate, but enter the lymph node paracortex to become interdigitating cells. Scand. J. Immunol. 27, 97.

Fossuam, S. (1989b) The life history of dendritic leukocytes [DL]. In: O.H. Ivessen (Ed.), Current Topics in Pathology. Springer-Verlag, Berlin, p. 101.

5 Hart, D.N.J. and McKenzie, J.L. (1990) Interstitial dendritic cells. Intl. Rev. Immunol. 6, 128.

Helfgott, D.C., Tatter, S.B., Santhanam, U., Clarick, R.H., Bhardwaj, N., May, L.T. and Sehgal, P.B. (1988) Interferon-beta 2/interleukin-6 in plasma and body fluids during acute bacterial infection. J. Immunol. 142, 948.

Inaba, K., Metlay, J.P., Crowley, M.T. and Steinman, R.M. (1990a) Dendritic cells pulsed with protein antigens in vitro can prime antigen-specific, MHC-restricted T cells in situ. J. Exp. Med. 172, 631.

Inaba, K., Metlay, J.P., Crowley, M.T., Witmer-Pack, M. and Steinman, R.M. (1990b) Dendritic cells as antigen presenting cells in vivo. Intl. Rev. Immunol. 6, 197.

20 Inaba, K., Steinman, R.M., Witmer-Pack, M., Aya, K., Inaba, M., Sudo, T., Wolpe, S. and Schuler, G. (1992) Identification of Proliferating Dendritic Cell Precursors In Mouse Blood. J. Exp. Med., 175:1157.

25 Jonuliet, H., Kohn, U., Muller, G., Wolfl, M., Lohmann, S., Saloga, J., Becker, D., Knop, J., and A.H. Enk, A.H. (1995). Keratinocyte-derived IL-15 enhances accessory functions of epidermal Langerhans cells and blood dendritic cells. J. Invest. Dermatol. 105, 861 [abstract].

30 Kashithara, M., Ueda, M., Horiguchi, Y., Furukawa, F., Hanaoka, M. and Imamura, S. (1986). A monoclonal antibody specifically reactive to human Langerhans cells. J. Invest. Dermatol. 87, 602.

-66 -

- Klareskog, L., Ronnelid, J. and Holm, G. (1995)  
Immunopathogenesis and immunotherapy in rheumatoid  
arthritis: an area in transition. J. Int. Med. 238, 191.
- Lanzavecchia, A. (1985). Antigen-specific  
interaction between T and B cells. Nature 314, 537.
- 5       Larsen, C.P., Ritchie, S.C., Hendrix, R., Linsley,  
P.S., Hathcock, K.S., Hodes, R.J., Lowry, R.P. and Pearson,  
T.C. (1994). Regulation of immunostimulatory function and  
costimulatory molecule (B7-1 and B7-2) expression on  
murine dendritic cells. J. Immunol. 152, 5208.
- 10       Lenz, A., Heine, M., Schuler, G. and Romani, N.  
(1993). Human and murine dermis contain dendritic cells.  
J. Clin. Invest. 92, 2587.
- 15       Mason, D.W., Pugh, C.W. and Webb, M. (1981) The rat  
mixed lymphocyte reaction: roles of a dendritic cell in  
intestinal lymph and T cell subsets defined by monoclonal  
antibodies. Immunol. 44, 75.
- 20       Mayordomo, J.I., Zorina, T., Storkus, W.J.,  
Zitvogel, L., Celluzzi, C., Falo, L.D., Melief, C.J.,  
Ilstad, S.T., Kast, W.M., DeLeo, A.B. and Lotze, M.T.  
(1995) Bone marrow-derived dendritic cells pulsed with  
synthetic tumour peptides elicit protective and  
therapeutic antitumor immunity. Nature Med. 1, 1297.
- 25       McWilliam, A.S., Nelson, D., Thomas, J.A. and Holt,  
P.G. (1994) Rapid dendritic cell recruitment is a hallmark  
of the acute inflammatory response at mucosal surfaces.  
J. Exp. Med. 179, 1331.
- 30       Mosialos, G., Birkenbach, M., Ayehunie, S.,  
Matsumura, F., Pinkus, G.S., Kieff, E. and Langhoff, E.  
(1996) Circulating human dendritic cells differentially  
express high levels of a 55-KD actin bundling protein. Am.  
J. Pathol. In Press.
- 35       O'Doherty, U., Steinman, R.M., Peng, M., Cameron,  
P.U., Gezelter, S., Kopeloff, I., Swiggard, W.J., Pope, M.  
and Bhardwaj, N. (1993) Dendritic cells freshly isolated

o from human blood express CD4 and mature into typical immunostimulatory dendritic cells after culture in monocyte-conditioned medium. J. Exp. Med. 178, 1067.

O'Doherty, U., Peng, M., Gezelter, S., Swiggard, W.J., Betjes, M., Bhardwaj, N. and Steinman, R.M. (1994) Human blood contains two subsets of dendritic cells, one  
5 immunologically mature, and the other immature. Immunol. 82, 487.

Pope, M., Betjes, M.G.H., Hirmand, H., Hoffman, L. and Steinman, R.M. (1995) Both dendritic cells and memory  
10 T lymphocytes emigrate from organ cultures of human skin and form distinctive dendritic T-cell conjugates. J. Invest. Dermatol. 104, 11.

Porgador, A. and Gilboa, E. (1995) Bone marrow-generated dendritic cells pulsed with a class  
15 I-restricted peptide are potent inducers of cytotoxic T lymphocytes. J. Exp. Med. 182, 255.

Pugh, C.W., MacPherson, G.G. and Steer, H.W. (1983) Characterization of nonlymphoid cells derived from rat peripheral lymph. J. Exp. Med. 157, 1758.

Rabinowitz, S.S. and Gordon, S. (1991) Macrosialin, a macrophage-restricted membrane sialoprotein  
20 differentially glycosylated in response to inflammatory stimuli. J. Exp. Med. 174, 827.

Ramprasad, M.P., Fischer, W., Witztum, J.L., Sambrano, G.R., Quehenberger, O. and Steinberg, D. (1995)  
25 The 94- to 97-kDa mouse macrophage membrane protein that recognizes oxidized low density lipoprotein and phosphatidylserine-rich liposomes is identical to macrosialin, the mouse homologue of human CD68. Proc. Natl. Acad. Sci. USA 92, 9580.

30 Reid, C.D.L., Stackpoole, A. Meager, A. and Tikerpae, J. (1992) Interactions of tumor necrosis factor with granulocyte-macrophage colony stimulating factor and other cytokines in the regulation of dendritic cell growth

- o in vitro from early biopotent CD34<sup>+</sup> progenitors in human bone marrow. J. Immunol. 149, 509.

Reis e Sousa, C., Stahl, P.D. and Austyn, J.M. (1993) Phagocytosis of antigens by Langerhans cells in vitro. J. Exp. Med. 178, 509.

- 5 Romani, N., Gruner, S., Brang, D., Kampgen, E., Lenz, A., Trockenbacher, B., Konwalinka, G., Fritsch, P.O., Steinman, R.M. and Schuler, G. (1994) Proliferating dendritic cell progenitors in human blood. J. Exp. Med. 180, 83.

- 10 Romani N., Koide, S., Crowley, M., Witmer-Pack, M., Livingstone, A.M., Fathman, C.G., Inaba, K. and Steinman, R.M. (1989a) Presentation of exogenous protein antigens by dendritic cells to T cell clones: intact protein is presented best by immature epidermal Langerhans cells. J. Exp. Med. 169, 1169.

- 15 Romani, N., Lenz, A., Glassel, H., Stossel, H., Stanzi, U., Majdic, O., Fritsch, P. and Schuler, G. (1989b) Cultured human Langerhans cells resemble lymphoid dendritic cells in phenotype and function. J. Invest. Dermatol. 93, 600.

- 20 Sallusto, F., Cella, M., Danieli, C. and Lanzavecchia, A. (1995) Dendritic cells use macropinocytosis and the mannose receptor to concentrate macromolecules in the major histocompatibility complex class II compartment: Downregulation by cytokines and bacterial products. J. Exp. Med. 182, 389.

- 25 Sallusto, F. and Lanzavecchia, A. (1994) Efficient presentation of soluble antigen by cultured human dendritic cells is maintained by granulocyte/macrophage colony-stimulating factor plus interleukin 4 and downregulated by tumor necrosis factor  $\alpha$ . J. Exp. Med. 30 179, 1109.

Sallusto, F. and Lanzavecchia, A. (1995) Dendritic cells use macropinocytosis and the mannose receptor to

- o concentrate antigen in the MHC class II compartment.  
Downregulation by cytokines and bacterial products. J.  
Exp. Med. 182, 389.

Schuler, G. and Steinman, R.M. (1985) Murine  
epidermal Langerhans cells mature into potent  
immunostimulatory dendritic cells in vitro. J. Exp. Med.  
5 161, 526.

Steinman, R.M. (1991) The dendritic cell system and  
its role in immunogenicity. Annu. Rev. Immunol. 9, 271.

Steinman, R.M., Hoffman, L. and Pope, M. (1995)  
Maturation and migration of cutaneous dendritic cells. J.  
10 Invest. Dermatol. 105 2S.

Teunissen, M.B.M., Wormmeester, J., Krieg, S.R.,  
Peters, P.J., Vogels, I.M.C., Kapsenberg, M.L., and Bos,  
J.D. (1990) Human epidermal Langerhans cells undergo  
profound morphological and phenotypical changes during in  
15 vitro culture. J. Invest. Dermatol. 94, 166.

Thomas, R., Davis, L.S. and Lipsky, P.E. (1993)  
Isolation and characterization of human peripheral blood  
dendritic cells. J. Immunol. 150, 821.

Thomas, R., Davis, L.S. and Lipsky, P.E. (1994)  
20 Rheumatoid synovium is enriched in mature  
antigen-presenting dendritic cells. J. Immunol. 152,  
2613.

Watson, S.R., Imai, Y., Fennie, C., Geoffroy, J.S.,  
Rosen, S.D. and Lasky, L.A. (1990) A homing receptor-IgG  
25 chimera as a probe for adhesive ligands of lymph node high  
endothelial venules. J. Cell Biol. 110, 2221.

Young, J.W. and Steinman, R.M. (1988) Accessory cell  
requirements for the mixed-leukocyte reaction and  
polyclonal mitogens, as studied with a new technique for  
30 enriching blood dendritic cells. Cell. Immunol. 111, 167.

Young, J.W. and Steinman, R.M. (1990) Dendritic  
cells stimulate primary human cytolytic lymphocyte

- o responses in the absence of CD4<sup>+</sup> helper T cells. J. Exp. Med. 171, 1315.

Zhou, L.-J., Schwarting, R., Smith, H.M. and Tedder, T.F. (1992) A novel cell-surface molecule expressed by human interdigitating reticulum cells, langerhans cells, and activated lymphocytes is a new member of the Ig  
5 superfamily. J. Immunol. 149, 735.

Zhou, L.-J. and Tedder, T.F. (1995) Human blood dendritic cells selectively express CD83, a member of the immunoglobulin superfamily. J. Immunol. 154, 3821.

Zitvogel, L., Mayordomo, J.I., Tjandrawan, T.,  
10 DeLeo, A.B., Clarke, M.R., Lotze, M.T. and Storkus, W.J. (1996) Therapy of murine tumors with tumor peptide pulsed dendritic cells: Dependence on T-cells, B7 costimulation, and Th1-associated cytokines. J. Exp. Med. In press.

Zvaifler, N.J., Steinman, R.M., Kaplan, G., Lau,  
15 L.L. and Rivelis, M. (1985) Identification of immunostimulatory dendritic cells in the synovial effusions of patients with rheumatoid arthritis. J. Clin. Invest. 76, 789.

20 All of the foregoing are incorporated herein by reference.

25

30

35



## WE CLAIM:

1. An in vitro method of producing mature dendritic cells from immature dendritic cells derived from pluripotential cells having the potential of expressing either macrophage or dendritic cell characteristics, said method comprising contacting the immature dendritic cells with a dendritic cell maturation factor for a time sufficient for said immature dendritic cells to mature and express dendritic cell characteristics.
2. The method according to claim 1 wherein the dendritic cell maturation factor is monocyte conditioned medium comprising said maturation factor.
3. The method according to claim 1 wherein said immature dendritic cells are derived from pluripotential cells cultured in the presence of at least one cytokine.
4. The method according to claim 3 wherein the cytokine is a combination of GM-CSF and either one or both of IL-4 and IL-13.
5. The method according to claim 4 wherein said pluripotential cells are human.
6. The method according to claim 5 wherein said pluripotential cells are obtained from human blood.
7. The method according to claim 6 wherein the cytokine is a combination of GM-CSF and IL-4.
8. A method of activating T cells comprising:
  - a) obtaining a population of pluripotential cells having the potential of expressing either macrophage

-72 -

- or dendritic cell characteristics from an individual and placing them in culture;
- b) contacting the pluripotential cells of step a with at least one cytokine to produce immature dendritic cells;
- 5 c) contacting the immature dendritic cells with a dendritic cell maturation factor for a time sufficient to cause said immature dendritic cells to express dendritic cell characteristics;
- 10 d) contacting the dendritic cells obtained from step c with an antigen to produce primed dendritic cells;
- e) exposing T cells to the primed dendritic cells of step d to activate said T cells.
9. The method according to claim 8 wherein the  
15 cytokine is a combination of GM-CSF and one or both of IL-4 or IL-13.
10. The method according to claim 9 wherein the  
20 dendritic cell maturation factor is monocyte conditioned medium.
11. The method according to claim 8 wherein the  
25 antigen is selected from the group consisting of fungal, viral, bacterial, tumor or self antigens.
12. The method according to claim 8 wherein the  
pluripotential cells are obtained from human blood.
13. The method according to claim 12 wherein the  
30 cytokine for contacting the pluripotential cells from human blood is a combination of GM-CSF and IL-4.

35

14. The method according to claim 8 wherein the T cells to be activated include antigen specific helper T cells.

15. The method according to claim 8 wherein the T cells to be activated include antigen specific killer T cells.

16. An assay to detect a dendritic cell maturation factor comprising:

a) contacting a test substance with a culture of unstable, immature dendritic cells derived from a population of pluripotential cells having the potential of expressing either macrophage or dendritic cell characteristics and which express characteristics of immature dendritic cells when cultured in a medium containing at least one cytokine;

b) detecting the maturation of the immature dendritic cells in response to the presence of the test substance.

17. The method according to claim 16 wherein maturation is detected by retention of a dendritic cell phenotype for three days following removal of the test substance from the culture medium contacting the dendritic cells.

18. The method according to claim 16 wherein maturation of the immature dendritic cells is detected by detecting the expression of an antigen selected from the group consisting of p55, CD83 and CD86.

19. The method according to claim 16 wherein maturation of the immature dendritic cells is detected by detecting a decrease in expression of CD115 antigen on the

- o mature dendritic cells compared to the immature dendritic cells.

20. A culture medium useful for causing the maturation of immature dendritic cells to phenotypically stable, mature dendritic cells, the culture medium  
5 comprising a mixture of salts, carbohydrates and vitamins at physiologic concentrations, about 1-5% human serum or plasma, GM-CSF and IL-4 in amounts together sufficient to promote maturation of pluripotential PBMC to immature  
10 dendritic cells, and a sufficient concentration of PBMC conditioned medium to cause the maturation of immature dendritic cells to stable mature dendritic cells.

21. The culture medium according to claim 20 wherein the human serum is present at a concentration of about 1-  
15 2%.

22. The culture medium according to claim 21 wherein the human serum is autologous to the source of immature  
20 dendritic cells.

23. A culture of mature dendritic cells prepared according to the method of claim 1.

24. The culture according to claim 23 wherein the  
25 cells are human and the dendritic cell maturation factor is monocyte conditioned medium.

25. A method of increasing the number of mature  
30 dendritic cells in an individual comprising administering to the individual a dendritic cell maturation factor in an amount sufficient to cause maturation of immature dendritic cells to mature dendritic cells.

26. The method according to claim 25 wherein the individual is pretreated with at least one cytokine to increase the endogenous number of immature dendritic cells.

5 27. The method according to claim 26 wherein the cytokine is selected from the group consisting of G-CSF, GM-CSF, IL-4 and IL-13.

10 28. The method according to claim 25 wherein the dendritic cell maturation factor is derived from PBMC conditioned medium.

15

20

25

30

35

1/14




FIG. 1

2/14




FIG. 2

3/14




FIG. 2



4/14



LINEAGE RESTRICTED ANTIGENS

FIG. 2

*FIG. 3*




FIG. 4





FIG. 5

*FIG. 6*




FIG. 7





FIG. 8



*FIG. 9*



12/14



*FIG. 10*

13/14




FIG. 11




FIG. 12

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 97/02110

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C12N5/08 C12Q1/02 A61K35/14 A61K38/19

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	THE JOURNAL OF INVESTIGATIVE DERMATOLOGY, vol. 105, no. 1, 1995, pages 25-75, XP000676459 STEINMAN R. ET AL: "Maturation and migration of cutaneous dendritic cells" see page 25 - page 35 ---	1-16, 20-28
X	JOURNAL OF EXPERIMENTAL MEDICINE, vol. 178, 1993, pages 1067-1078, XP000676564 O'DOHERTY U. ET AL: "Dendritic cells freshly isolated from human blood express CD4 and mature into typical immunostimulatory dendritic cells after culture in monocyte-conditioned medium" see the whole document --- -/--	1-7, 16, 23-25

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*A\* document member of the same patent family

Date of the actual completion of the international search

1 July 1997

Date of mailing of the international search report

15. 07. 97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Fernandez y Branas, F

# INTERNATIONAL SEARCH REPORT

nal Application No

PCT/US 97/02110

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>IMMUNOLOGY, vol. 82, 1994, pages 487-493, XP000676573 O'DOHERTY U. ET AL: "Human blood contains two subsets of dendritic cells, one immunologically mature and the other immature" see the whole document ---</p>	1-7, 16, 23-25
X,P	<p>JOURNAL OF IMMUNOLOGICAL METHODS, vol. 196, 1996, pages 121-135, XP002034181 BENDER, A. ET AL: "Improved methods for the generation of dendritic cells from nonproliferating progenitors in human blood" see the whole document ---</p>	1-28
X,P	<p>JOURNAL OF IMMUNOLOGICAL METHODS, vol. 196, 1996, pages 137-151, XP002034182 ROMANI N. ET AL: "Generation of mature dendritic cells from human blood. An improved method with special regard to clinical applicability" see the whole document ---</p>	1-28
A	<p>JOURNAL OF EXPERIMENTAL MEDICINE, vol. 180, 1994, pages 83-93, XP002034183 ROMANI, N. ET AL: "Proliferating dendritic cell progenitors in human blood" see the whole document ---</p>	1-28
A	<p>JOURNAL OF EXPERIMENTAL MEDICINE, vol. 179, 1994 - 1109, page 1118 XP000676465 SALLUSTO F. ET AL: "Efficient presentation of soluble antigen by cultured human dendritic cells is maintained by granulocyte/macrophage colony-stimulating factor plus interleukin-4 and downregulated by tumor necrosis factor-alpha" see the whole document ---</p>	1-28
A	<p>THE JOURNAL OF IMMUNOLOGY, vol. 154, 1995, pages 3821-3835, XP002034184 ZHOU L-J. ET AL: "Human dendritic cells selectively express CD83, a member of the immunoglobulin superfamily" see the whole document ---</p>	1-28

-/--

# INTERNATIONAL SEARCH REPORT

Inter. Application No

PCT/US 97/02110

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>NATURE MEDICINE, vol. 2, no. 1, January 1996, pages 52-58, XP000605401 HSU, F.J. ET AL: "Vaccination of patients with B-cell lymphoma using autologous antigen-pulsed dendritic cells" see the whole document ---</p>	1-28
A	<p>WO 95 28479 A (KLINIKUM DER ALBERT LUDWIGS UN ;KANZ LOTHAR (DE); BRUGGER WOLFRAM) 26 October 1995 see the whole document -----</p>	1-28

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 97/02110

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claim(s) 25-28  
is(are) directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Inte Application No

PCT/US 97/02110

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9528479 A	26-10-95	DE 4412794 A	14-12-95
		AU 2302495 A	10-11-95
		CA 2187770 A	26-10-95
		EP 0755439 A	29-01-97
-----			